



- 1 -

TITLE OF THE INVENTIONMELANOMA ANTIGENS AND THEIR USE IN  
DIAGNOSTIC AND THERAPEUTIC METHODS.

5

FIELD OF THE INVENTION

This invention is in the field of prevention and treatment of human cancers. More specifically, this invention relates to genes encoding melanoma antigens recognized by T-Cells and their corresponding proteins and to preventative, diagnostic and therapeutic applications which employ these genes or proteins.

BACKGROUND OF THE INVENTION

Melanomas are aggressive, frequently metastatic tumors derived from either melanocytes or melanocyte related nevus cells ("Cellular and Molecular Immunology" (1991) (eds) Abbas A.K., Lechtman, A.H., Pober, J.S.; W.B. Saunders Company, Philadelphia: pages 340-341). Melanomas make up approximately three percent of all skin cancers and the worldwide increase in melanoma is unsurpassed by any other neoplasm with the exception of lung cancer in women ("Cellular and Molecular Immunology" (1991) (eds) Abbas, A.K., Lechtman, A.H., Pober, J.S.; W.B. Saunders Company Philadelphia pages: 340-342; Kirkwood and Agarwala (1993) Principles and Practice of Oncology 7:1-16). Even when melanoma is apparently localized to the skin, up to 30% of the patients will develop systemic metastasis and the majority will die (Kirkwood and Agarwala (1993) Principles and Practice of Oncology 7:1-16). Classic modalities of treating melanoma include surgery, radiation and chemotherapy. In the past decade immunotherapy and gene therapy have emerged as new and promising methods for treating melanoma.

T cells play an important role in tumor regression in most murine tumor models. Tumor infiltrating lymphocytes (TIL) that recognize unique cancer antigens can be isolated from many murine tumors. The adoptive transfer of

these TIL plus interleukin-2 can mediate the regression of established lung and liver metastases (Rosenberg, S.A., et al., (1986) Science 233:1318-1321). In addition, the secretion of IFN- $\gamma$  by injected TIL significantly correlates with in vivo regression of murine tumors suggesting activation of T-cells by the tumor antigens. (Barth, R.J., et al., (1991) J. Exp. Med. 173:647-658). The known ability of tumor TIL to mediate the regression of metastatic cancer in 35 to 40% of melanoma patients when adoptively transferred into patients with metastatic melanoma attests to the clinical importance of the antigens recognized (Rosenberg, S.A., et al., (1988) N Engl J Med 319:1676-1680; Rosenberg S.A. (1992) J. Clin. Oncol. 10:180-199).

T cell receptors on CD8<sup>+</sup> T cells recognize a complex consisting of an antigenic peptide (9-10 amino acids for HLA-A2),  $\beta$ -2 microglobulin and class I major histocompatibility complex (MHC) heavy chain (HLA-A, B, C, in humans). Peptides generated by digestion of endogenously synthesized proteins are transported into the endoplasmic reticulum, bound to class I MHC heavy chain and  $\beta$ 2 microglobulin, and finally expressed in the cell surface in the groove of the class I MHC molecule. Therefore, T cells can detect molecules that originate from proteins inside cells, in contrast to antibodies that detect intact molecules expressed on the cell surface. Therefore, antigens recognized by T cells may be more useful than antigens recognized by antibodies.

Strong evidence that an immune response to cancer exists in humans is provided by the existence of lymphocytes within melanoma deposits. These lymphocytes, when isolated, are capable of recognizing specific tumor antigens on autologous and allogeneic melanomas in an MHC restricted fashion. (Itoh, K. et al. (1986), Cancer Res. 46: 3011-3017; Muul, L.M., et al. (1987), J. Immunol. 138:989-995); Topalian, S.L., et al., (1989) J. Immunol. 142: 3714-3725; Darrow, T.L., et al., (1989) J. Immunol.

142: 3329-3335; Hom, S.S., et al., (1991) J. Immunother. 10:153-164; Kawakami, Y., et al., (1992) J. Immunol. 148: 638-643; Hom, S.S., et al., (1993) J. Immunother. 13:18-30; O'Neil, B.H., et al., (1993) J. Immunol. 151: 1410-1418). TIL from patients with metastatic melanoma  
5 recognize shared antigens including melanocyte-melanoma lineage specific tissue antigens in vitro (Kawakami, Y., et al., (1993) J. Immunother. 14: 88-93; Anichini, A. et al., (1993) et al., J. Exp. Med. 177: 989-998). Anti-melanoma T cells appear to be enriched in TIL probably as  
10 a consequence of clonal expansion and accumulation at the tumor site in vivo (Sensi, M., et al., (1993) J. Exp. Med. 178:1231-1246). The fact that many melanoma patients mount cellular and humoral responses against these tumors and that melanomas express both MHC antigens and tumor  
15 associated antigens (TAA) suggests that identification and characterization of additional melanoma antigens will be important for immunotherapy of patients with melanoma.

Peripheral blood lymphocytes have been used to identify potential melanoma tumor antigens. Van Der  
20 Bruggen et al. (1991) Science 254: 1643-1647 has characterized a gene coding for a melanoma antigen, designated MAGE-1, using T cell clones established from the peripheral blood of patients who were repetitively  
25 immunized in vivo with mutagenized tumor cells. Cytotoxic T-cells derived from the peripheral blood lymphocytes of patients with melanoma were used to identify a potential antigenic peptide encoding MAGE-1 (Traversari, C., et al. (1992) J. Exp. Med. 176:1453-1457). Brichard et al. (1993) J. Exp. Med. 178:489-495 has also characterized a  
30 gene encoding a melanoma antigen designated tyrosinase using peripheral blood lymphocytes from patients who were sensitized by repetitive in vitro stimulation with tumor. Further support for the therapeutic potential of melanoma antigens is provided by Brown et al. (United States Patent  
35 No. 5,262,177). Brown et al. (United States Patent Number 5,262,177) relates to a recombinant vaccinia virus-based

4

° melanoma vaccine where the melanoma antigen p97 is and reported to show a protective effect from tumor cell challenge in both murine model. Characterization of additional melanoma antigens be important for the development of new strategies for cancer immunotherapy, in particular for melanoma.

#### SUMMARY OF THE INVENTION

This invention relates, in general, to nucleic acid sequences encoding melanoma antigens recognized by T-lymphocytes (MART-1) and protein and peptide encoded by these sequences. This invention further provides bioassays for these nucleic acid sequences, proteins and peptides. This invention also provides therapeutic uses for the nucleic acid sequences, proteins or peptides described herein.

It is a general of the present invention to provide a substantially purified and isolated nucleic acid sequence which encodes for the MART-1 melanoma antigen.

It is another object of this invention to provide a recombinant molecule comprising a vector and all or part of the nucleic acid sequence encoding MART-1.

It is another object of this invention to produce recombinant proteins encoded by all or part of the nucleic acid sequence encoding MART-1.

It is a further object of this invention to provide monoclonal or polyclonal antibodies reactive with the MART-1 protein, peptides or portions thereof.

It is an object of this invention to provide methods of detecting the MART-1 gene or MART-1 mRNA in a biological sample.

It is another object of this invention to provide methods of detecting the MART-1 protein or peptides in a biological sample.

It is an object of this invention to provide diagnostic methods for human disease, in particular for melanomas and metastatic melanomas.

It is a further object of this invention to provide

5

° methods for prophylactic or therapeutic uses involving all or part of the nucleic acid sequence encoding MART-1 and its corresponding protein or peptides derived from the MART-1 amino acid sequence.

5 It is also an object of this invention to provide melanoma vaccines comprising all or part of the nucleic acid sequence encoding MART-1 or its corresponding protein for preventing or treating melanoma.

10 It is a further object of this invention to provide immunogenic peptides derived from the MART-1 protein sequence for use in vaccines.

In addition, it is another object of this invention to provide multivalent vaccines comprising all or part of the MART-1 nucleic acid sequence or its corresponding protein or peptides and at least one other immunogenic molecule capable of eliciting the production of antibodies in a mammal to melanoma antigens.

20 It is another object of this invention to provide a method for preventing or treating melanoma utilizing all or part of the MART-1 nucleic acid sequence or its corresponding protein in gene therapy protocols.

It is a further object of this invention to provide immunogenic peptides derived from a gp100 melanoma antigen protein sequence for use in vaccines.

25 It is yet another object of this invention to provide a method of prophylactic or therapeutic immunization for melanoma using the vaccines described herein.

It is a further object of this invention to provide a method of identifying melanoma antigens that would constitute potential targets for immunotherapy.

30 It is yet another object of this invention to provide a method of identifying candidate immunogenic peptides derived from either the MART-1 or gp100 sequences for use in immunotherapy.

35

#### DESCRIPTION OF THE FIGURES

Figure 1 shows the nucleotide and predicted amino

6

acid sequence of the cDNA encoding the MART-1 antigen. The hydrophobic region is underlined.

Figure 2 shows titration of MART-1 peptides for recognition by TIL. T2 cells were incubated with varied concentrations of the purified MART-1 peptides, M9-1, M9-2, M9-3, M10-2, M10-3, M10-4 and M10-5, and lysis by TIL clone A42 or TIL line TIL1235 was measured by 4h-<sup>51</sup>Cr release cytotoxicity assay at an E (EFFECTOR):T (TARGET) ratio of 20:1 for A42 and 40:1 for TIL1235. Peptide M9-2 sensitized T2 cells at a concentration of 1ng/ml. The purified peptide M10-4 was recognized by TIL1235, but not by A42. (M9-1 | - |, M9-2 ⊙-⊙, M9-3 ⊠-⊠, M10-2 Δ-Δ, M10-3 ▽-▽, M10-4 ⊞-⊞, M10-5 +-+ ).

Figure 3A shows a radionuclide scan of patient 1200 with metastatic melanoma after receiving the adoptive transfer of autologous <sup>111</sup>In labeled TIL1200. The arrow indicates one of the areas of TIL accumulation corresponding to a metastatic lesion in the left thigh.

Figure 3B shows regression of subcutaneous metastatic tumors following treatment with TIL1200 plus IL-2.

Treatment began on day 0.

Figure 4 show the nucleic acid sequence of the full length cDNA25. The start and stop codons are underlined.

Figure 5A shows amino acid sequence of the full length cDNA25. The antigenic peptide is underlined.

Figure 5B shows comparison of the amino acid sequence of the full length cDNA25 (cDNA25FL), the truncated form of cDNA25 (cDNA25TR), Pmel17, ME20 and gp100. (⊙ indicates deletion; - indicates identity).

Figure 6 shows northern blot analysis of melanoma and neonatal melanocyte cell lines and various fresh tissues (10-20ug of total RNA) with a cDNA25 probe (the Sal I digested fragment of pCRII-cDNA25) and the β-actin probe (Clontech). C32, 586mel melanoma cell lines and NHEM529, NHEM530 neonatal melanocyte cell lines were very weak positive.

#### DETAILED DESCRIPTION OF THE INVENTION

7

For the purpose of a more complete understanding of the invention, the following definitions are described herein. Nucleic acid sequences includes, but is not limited to, DNA, RNA or cDNA. Nucleic acid sequence as used herein refers to an isolated and purified nucleic acid sequence. MART-1 messenger RNA (mRNA) refers to one or more RNA transcripts which are a product of the MART-1 gene. Substantially homologous as used herein refers to substantial correspondence between the nucleic acid sequence of MART-1 shown in Figure 1 (SEQ ID NO:1) and that of any other nucleic acid sequence. Substantially homologous means about 50-100% homologous homology, preferably by about 70-100% homology, and most preferably about 90-100% homology between the MART-1 sequence and that of any other nucleic acid sequence. In addition, substantially homologous as used herein also refers to substantial correspondences between the amino acid sequence of the MART-1 antigen shown in Figure 1 (SEQ ID NO:2) and that of any other amino acid sequence.

Major Histocompatibility Complex (MHC) is a generic designation meant to encompass the histo-compatibility antigen systems described in different species including the human leucocyte antigens (HLA).

The term melanoma includes, but is not limited to, melanomas, metastatic melanomas, melanomas derived from either melanocytes or melanocytes related nevus cells, melanocarcinomas, melanoepitheliomas, melanosarcomas, melanoma in situ, superficial spreading melanoma, nodular melanoma, lentigo maligna melanoma, acral lentiginous melanoma, invasive melanoma or familial atypical mole and melanoma (FAM-M) syndrome. Such melanomas in mammals may be caused by, chromosomal abnormalities, degenerative growth and developmental disorders, mitogenic agents, ultraviolet radiation (UV), viral infections, inappropriate tissue expression of a gene, alterations in expression of a gene, and presentation on a cell, or carcinogenic agents. The aforementioned melanomas can be

° diagnosed, assessed or treated by methods described in the present application.

By atypical mole we mean a mole with features that are abnormal and may be precancerous.

5 By melanoma antigen or immunogen we mean all or parts thereof of the MART-1 protein or peptides based on the MART-1 protein sequence capable of causing a cellular or humoral immune response in a mammal. Such antigens are also reactive with antibodies from animals immunized with all, part or parts of the MART-1 protein (SEQ ID NO:2).  
10 Such a protein or peptide may be encoded by all or part of the MART-1 nucleic acid sequence of this invention.

By immunogenic peptide we mean a peptide derived from the MART-1 protein sequence (FIGURE 1; SEQ ID NO. 2) or a gp100 protein sequence (FIGURE 5A; SEQ ID NO: 27) capable  
15 of causing a cellular or humoral immune response in a mammal. Such peptides may also be reactive with antibodies from an animal immunized with the peptides. Such peptides are about 5-20 amino acid in length preferably about 8 to 15 amino acids in length, and most  
20 preferably about 9-10 amino acids in length.

One skilled in the art will understand that the bioassays of the present invention may be used in the analysis of biological samples or tissues from any vertebrate species. In a preferred embodiment, mammalian  
25 biological samples or tissues are analyzed.

Tissue includes, but is not limited to, single cells, whole organs and portions thereof. Biological samples include, but are not limited to, tissues, primary cultures of mammalian tissues, biopsy specimens, pathology  
30 specimens, and necropsy specimens. Mammal includes but is not limited to, humans, monkeys, dogs, cats, mice, rats, pigs, cows, pigs, horses, sheep and goats.

The present invention provides a nucleic acid sequence which encodes a novel melanoma antigen recognized  
35 by T cells. This novel melanoma antigen designated MART-1 (melanoma antigen recognized by T-Cells-1). MART-1 shows



no significant homology to any known melanoma antigen and thus represents a new melanoma antigen. The MART-1 antigen contains a highly hydrophobic region from amino acids 27 to 47 (SEQ ID. NO:2) followed by three arginine residues, suggestive of a transmembrane protein. Although no significant homology exists to the entire protein there is a 27 amino acid segment (amino acids 57-83; SEQ ID. NO:2) that is 37% identical to a Type II membrane protein previously recognized as mouse natural killer cell surface protein NKR-P1 (Yokoyama, W.M., et al. (1991), J. Immunol. 147:3229-3236). MART-1 does not contain a leader sequence characteristic of many Type I membrane proteins (Singer, S.J. (1990) Annu. Rev. Cell Biol. 6: 247-296).

MART-1 RNA expression appears to be restricted to fresh and cultured melanoma and melanocyte cell lines and human retina; expression has not been found in any other fresh or cultured tissues or other tumor histologies tested. The cDNA sequence for MART-1 is shown in Figure 1 (SEQ ID NO:1), the deduced amino acid sequence for the MART-1 protein is also shown in Figure 1 (SEQ ID NO.:1).

The nucleic acid sequence for MART-1 shown in Figure 1 (SEQ ID NO.:1), represents a preferred embodiment of the invention. It is, however, understood by one skilled in the art that due to the degeneracy of the genetic code variations in the cDNA sequence shown in Figure 1 (SEQ ID NO.:1) will still result in a DNA sequence capable of encoding the MART-1 protein antigen. Such DNA sequences are therefore functionally equivalent to the sequence set forth in Figure 1 (SEQ ID NO.:1) and are intended to be encompassed within the present invention. Further, a person of skill in the art will understand that there are naturally occurring allelic variations in a given species of the MART-1 nucleic acid sequence shown in Figure 1 (SEQ ID NO.:1), these variations are also intended to be encompassed by the present invention.

The predicted MART-1 antigen is a 118 amino acid protein of about 13 (kd). This invention further includes

° MART-1 protein or peptides or analogs thereof having substantially the same function as the MART-1 antigen or protein of this invention. Such proteins or polypeptides include, but are not limited to, a fragment of the protein, or a substitution, addition or deletion mutant of  
5 the MART-1 protein. This invention also encompasses proteins or peptides that are substantially homologous to the MART-1 antigen. Substantially homologous means about 50-100% homology, preferably about 70-100% homology, and most preferably about 90-100% homology between the MART-1  
10 and any another amino acid sequence or protein or peptide.

The term "analog" includes any polypeptide having an amino acid residue sequence substantially identical to the MART-1 sequence specifically shown herein (Figure 1; SEQ ID NO: 1) in which one or more residues have been  
15 conservatively substituted with a functionally similar residue and which displays the functional aspects of the MART-1 antigen as described herein. Examples of conservative substitutions include the substitution of one non-polar (hydrophobic) residue such as isoleucine,  
20 valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine  
25 or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid or another.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a  
30 non-derivatized residue. "Chemical derivative" refers to a subject polypeptide having one or more residues chemically derivatized by reaction of a functional side group. Examples of such derivatized molecules include for example, those molecules in which free amino groups have  
35 been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl

11

groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine  
5 may be derivatized to form N-im-benzylhistidine. Also included as chemical derivatives are those proteins or peptides which contain one or more naturally-occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for  
10 proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine. Proteins or polypeptides of the present invention also include any polypeptide having  
15 one or more additions and/or deletions or residues relative to the sequence of a polypeptide whose sequence is encoded is the DNA of MART-1, so long as the requisite activity is maintained.

This invention also provides a recombinant DNA  
20 molecule comprising all or part of the MART-1 nucleic acid sequence (SEQ. ID NO.:1) and a vector. Expression vectors suitable for use in the present invention comprise at least one expression control element operationally linked to the nucleic acid sequence. The expression control  
25 elements are inserted in the vector to control and regulate the expression of the nucleic acid sequence. Examples of expression control elements includes, but is not limited to, lac system, operator and promoter regions of phage lambda, yeast promoters and promoters derived  
30 from polyoma, adenovirus, retrovirus or SV40. Additional preferred or required operational elements include, but are not limited to, leader sequence, termination codons, polyadenylation signals and any other sequences necessary or preferred for the appropriate transcription and  
35 subsequent translation of the nucleic acid sequence in the host system. It will be understood by one skilled in the

art the correct combination of required or preferred expression control elements will depend on the host system chosen. It will further be understood that the expression vector should contain additional elements necessary for the transfer and subsequent replication of the expression vector containing the nucleic acid sequence in the host system. Examples of such elements include, but are not limited to, origins of replication and selectable markers. It will further be understood by one skilled in the art that such vectors are easily constructed using conventional methods (Ausubel et al., (1987) in "Current Protocols in Molecular Biology", John Wiley and Sons, New York, New York) or commercially available.

Another aspect of this invention relates to a host organism into which recombinant expression vector containing all or part of the MART-1 nucleic acid sequence has been inserted. The host cells transformed with the MART-1 nucleic acid sequence of this invention includes eukaryotes, such as animal, plant, insect and yeast cells and prokaryotes, such as E. coli. The means by which the vector carrying the gene may be introduced into the cell includes, but is not limited to, microinjection, electroporation, transduction, or transfection using DEAE-dextran, lipofection, calcium phosphate or other procedures known to one skilled in the art (Sambrook et al. (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York).

In a preferred embodiment, eukaryotic expression vectors that function in eukaryotic cells are used. Examples of such vectors include, but are not limited to, retroviral vectors, vaccinia virus vectors, adenovirus vectors, herpes virus vector, fowl pox virus vector, bacterial expression vectors, plasmids, such as pcDNA3 (Invitrogen, San Diego, CA) or the baculovirus transfer vectors. Preferred eukaryotic cell lines include, but are not limited to, COS cells, CHO cells, HeLa cells, NIH/3T3 cells, 293 cells (ATCC# CRL1573), T2 cells, dendritic

cells, or monocytes. In a particularly preferred embodiment the recombinant MART-1 protein expression vector is introduced into mammalian cells, such as NIH/3T3, COS, CHO, 293 cells (ATCC #CRL 1573), T2 cells, dendritic cells, or monocytes to ensure proper processing and modification of the MART-1 protein. In an alternative embodiment the MART-1 cDNA is introduced into COS7 (Gluzman, Y. et al. (1981) Cell 23: 175-182).

In one embodiment the expressed recombinant MART-1 protein may be detected by methods known in the art which include Coomassie blue staining and Western blotting using antibodies specific for the MART-1 protein.

In a further embodiment, the recombinant protein expressed by the host cells can be obtained as a crude lysate or can be purified by standard protein purification procedures known in the art which may include differential precipitation, molecular sieve chromatography, ion-exchange chromatography, isoelectric focusing, gel electrophoresis, affinity, and immunoaffinity chromatography and the like. (Ausubel et. al., (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New York). In the case of immunoaffinity chromatography, the recombinant protein may be purified by passage through a column containing a resin which has bound thereto antibodies specific for the MART-1 protein (Ausubel et. al., (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New York).

The nucleic acid sequence or portions thereof, of this invention are useful as probes for the detection of expression of the MART-1 gene in normal and diseased tissue. Therefore, another aspect of the present invention relates to a bioassay for detecting messenger RNA encoding the MART-1 protein in a biological sample comprising the steps of (a) contacting all or part of the nucleic acid sequence of this invention with said biological sample under conditions allowing a complex to

form between said nucleic acid sequence and said messenger RNA, (b) detecting said complexes and, (c) determining the level of said messenger RNA.

RNA can be isolated as whole cell RNA or as poly(A)<sup>+</sup> RNA. Whole cell RNA can be isolated by a variety of methods known to those skilled in the art. (Ausubel et al., (1987) on "Current Protocols in Molecular Biology", John Wiley and Sons, New York). Such methods include extraction of RNA by differential precipitation (Birnboim, H.C. (1988) Nucleic Acids Res., 16:1487-1497), extraction of RNA by organic solvents (Chomczynski, P. et al. (1987) Anal. Biochem., 162:156-159) and the extraction of RNA with strong denaturants (Chirgwin, J.M. et al. (1979) Biochemistry, 18:5294-5299). Poly(A)<sup>+</sup> RNA can be selected from whole cell RNA by affinity chromatography on oligo-d(T) columns (Aviv, H. et al. (1972) Proc. Natl. Acad. Sci., 69:1408-1412). Examples of methods for determining cellular messenger mRNA levels for step (c) include, but is not limited to Northern blotting (Alwine, J.C. et al. (1977) Proc. Natl. Acad. Sci., 74:5350-5354), dot and slot hybridization (Kafatos, F.C. et al. (1979) Nucleic Acids Res., 7:1541-1522), filter hybridization (Hollander, M.C. et al. (1990) Biotechniques; 9:174-179), RNase protection (Sambrook et. al., (1989) in "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Press, Plainview, NY), polymerase chain reaction (Watson, J.D. et al. (1992) in "Recombinant DNA" Second Edition, W.H. Freeman and Company, New York) and nuclear run-off assays (Ausubel et. al., (1987) in "Current Protocols in Molecular Biology" Supplement 9 (1990); John Wiley and Sons, New York, New York).

Detection of complexes in Step (b) of the bioassay can also be carried out by a variety of techniques. Detection of the complexes by signal amplification can be achieved by several conventional labelling techniques including radiolabels and enzymes (Sambrook et. al., (1989) in "Molecular Cloning, A Laboratory Manual", Cold

15

Spring Harbor Press, Plainview, New York; Ausubel et al., (1987) in "Current Protocols in Molecular Biology, John Wiley and Sons, New York New York). Radiolabelling kits are also commercially available. The MART-1 nucleic acid sequence used as a probe in step(c) of the bioassay may be RNA or DNA. Preferred methods of labelling the DNA sequences are with  $^{32}\text{P}$  using Klenow enzyme or polynucleotide kinase. Preferred methods of labeling RNA or riboprobe sequences are with  $^{32}\text{P}$  or  $^{35}\text{S}$  using RNA polymerases. In addition, there are known non-radioactive techniques for signal amplification including methods for attaching chemical moieties to pyrimidine and purine rings (Dale, R.N.K. et al. (1973) Proc. Natl. Acad. Sci., 70:2238-2242; Heck, R.F. (1968) S. Am. Chem. Soc., 90:5518-5523), methods which allow detection by chemiluminescence (Barton, S.K. et al. (1992) J. Am. Chem. Soc., 114:8736-8740) and methods utilizing biotinylated nucleic acid probes (Johnson, T.K. et al. (1983) Anal. Biochem., 133:125-131; Erickson, P.F. et al. (1982) J. of Immunology Methods, 51:241-249; Matthaei, F.S. et al (1986) Anal. Biochem., 157:123-128) and methods which allow detection by fluorescence using commercially available products. Non-radioactive labelling kits are also commercially available.

Examples of biological samples that can be used in this bioassay include, but are not limited to, primary mammalian cultures, continuous mammalian cell lines, such as melanocyte cell lines, mammalian organs such as skin or retina, tissues, biopsy specimens, neoplasms, pathology specimens, and necropsy specimens.

In a preferred embodiment, a  $^{32}\text{P}$  radiolabelled MART-1 probe, as exemplified in Example 1, is used. Preferably the MART-1 probe is the full length cDNA in Figure 1 (SEQ ID NO:1). The approximately 1.6 Kilobase (kb) cDNA (Figure 1; SEQ ID NO:1) was cloned into the vector and the resulting plasmid, deposited with the American Type Culture Collection (ATCC) 12301 Parklawn Drive, Rockville,

° MD 20852 USA on April 14, 1994, and given ATCC Deposit Number 75738. The full length MART-1 nucleic acid sequence can be isolated from the pCRII plasmid by digestion with HINDIII and XhoI restriction enzymes. This 1.6kb nucleic acid sequence can then be used as a probe. 5 This probe is used to detect MART-1 mRNA in total RNA or poly A<sup>+</sup> RNA isolated from a variety of tissues or biological samples.

In another embodiment, combinations of oligonucleotide pairs based on the MART-1 sequence in 10 Figure 1 (SEQ ID NO.:1) are used as Polymerase Chain Reaction (PCR) primers to detect MART-1 mRNA in a biological sample. These primers can be used in a method following the reverse transcriptase - Polymerase Chain Reaction (RT-PCR) process for amplifying selected RNA 15 nucleic acid sequences as detailed in Ausubel et al., (eds) (1987) In "Current Protocols in Molecular Biology" Chapter 15, John Wiley and Sons, New York, New York. The oligonucleotides can be synthesized by automated instruments sold by a variety of manufacturers or can be 20 commercially prepared based upon the nucleic acid sequence of this invention. One skilled in the art will know how to select PCR primers based on the MART-1 nucleic acid sequence for amplifying MART-1 RNA in a sample.

The MART-1 nucleic acid sequence or portions thereof 25 (Figure 1: SEQ ID NO:1) of this invention are useful to detect alterations of the MART-1 gene in normal or diseased mammalian tissue. By alteration, we mean additions, deletions, substitutions or duplications in the MART-1 gene sequence or gene amplification of the MART-1 30 gene sequence. Therefore, another aspect of the present invention relates to an assay for detecting alterations of the MART-1 gene in a biological sample comprising the steps of (a) contacting all or part of the nucleic acid sequence of this invention with genomic DNA isolated from 35 a biological sample under conditions allowing a complex to form between said nucleic acid sequence and said genomic



DNA, (b) detecting said complexes, and (c) determining alterations in said MART-1 gene by comparison to a control sample.

Standard methods for isolating DNA from a biological sample, detecting alterations in a gene and detecting complex between the MART-1 nucleic acid probe and genomic DNA sequences are provided in manuals such as Sambrook et al., (eds) (1989) "Molecular Cloning, A Laboratory Mineral", Cold Spring Harbor Press, Plainview, New York and in Ausubel et al., (eds) (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New York.

The MART-1 nucleic acid sequences of this invention (Figure 1; SEQ ID No:1) can also be used as probes to isolate the MART-1 homologs in other species. In a preferred embodiment the MART-1 cDNA (Figure 1; SEQ ID No:1) is used to screen a mammalian cDNA library, positive clones are selected and sequenced. Examples of tissue sources from which the cDNA library can be synthesized includes, but are not limited to skin, retina, melanocytes, neonatal skin and embryos. Preferably a melanoma library is screened using the MART-1 cDNA as a probe (Figure 1; SEQ ID No. 1). One skilled in the art will understand the appropriate hybridization conditions to be used to detect the homologs. Conventional methods for nucleic acid hybridization, construction of libraries and cloning techniques are described in Sambrook et al., (eds) (1989) In "Molecular Cloning A Laboratory Manual" Cold Spring Harbor Press, Plainview, New York and Ausubel et al., (eds) in "Current Protocols in Molecular Biology" (1987), John Wiley and Sons, New York, New York.

It is known that all or parts thereof of the MART-1 protein is an antigen present on melanoma cells. It is therefore another aspect of this invention to provide MART-1 nucleic acid probes to be utilized in detecting MART-1 RNA or alterations in the level of MART-1 mRNA in biological sample isolated from a mammal afflicted with a

° disease. Examples of such diseases, include but is not limited to, melanomas. By alterations in the level of MART-1 mRNA we mean an increase or decrease in the level of an RNA relative to a control sample or the appearance or disappearance of the MART-1 mRNA relative to a control sample. Detection in the alterations of MART-1 mRNA will allow for diagnosis or the assessment of the diseased state. Therefore, alterations in the level of MART-1 mRNA may be predictive of the prognosis for the afflicted mammal.

10 In another embodiment the nucleic acid of this invention can be used in in situ hybridization on mammalian tissues to determine the precise site or subcellular site of expression of the MART-1 gene within a tissue. A preferred method of labeling the MART-1 nucleic acid sequence is synthesizing a <sup>35</sup>S - labeled RNA probe by in vitro transcription utilizing SP6 polymerase. In the MART-1 plasmid (ATCC Deposit #75738) the sense strand is under the control of the T7 promoter, the antisense strand is under the SP6 promoter. It is preferable that the probe be hydrolyzed to a probe length of approximately 400-200 base pairs. Conventional methods for preparation of tissues for *in situ*, synthesis of probes and detection of signal can be found in Ausubel et. al., (eds) (1987) in "Current Protocols in Molecular Biology," John Wiley and Sons, New York, New York Chapter 14 and Vander Ploeg, M., Raap A.K. (1988) In "New Frontiers in Cytology" Goerttler, K., Feichter, GE, Witte. S. (eds) pp 13-21 Springer-Verlag, New York. The probe is then contacted with mammalian tissue sections and in situ analyses performed by conventional methods. Examples of tissues that can be used include, but are not limited to, mammalian embryos, adult mammalian tissues, such as skin, lymph nodes and retina, biopsy specimens, pathology specimens and necropsy specimens. In a preferred embodiment, MART-1 in situ probes may be used to evaluate MART-1 RNA expression in diseased tissue for invasive early melanoma to

° characterize radial and vertical growth phases of the melanoma lesion and assess the margins of the disease within the tissue.

In yet another embodiment of this invention all or parts thereof of the MART-1 (SEQ ID NO:1) nucleic acid  
5 sequence can be used to generate transgenic animals. Preferably the MART-1 gene is introduced into an animal or an ancestor of the animal at an embryonic stage, preferably at the one cell stage and generally not later than about the eight cell stage. There are several means  
10 by which transgenic animals carrying a MART-1 gene can be made. One method involves the use of retroviruses carrying all or part of the MART-1 sequence. The retroviruses containing the transgene are introduced into the embryonic animal by transfection. Another methods  
15 involves directly injecting the transgene into the embryo. Yet another methods employs the embryonic stem cell method or homologous recombination method known to workers in the field. Examples of animals into which the MART-1 transgene can be introduced includes but is not limited  
20 to, primates, mice, rats or other rodents. Such transgenic animals may be useful as biological models for the study of melanoma and to evaluate diagnostic or therapeutic methods for melanoma.

This invention further comprises an antibody or  
25 antibodies reactive with the MART-1 protein or peptides having the amino acid sequence defined in Figure 1 (SEQ ID NO: 2) or a unique portion thereof. In this embodiment of the invention the antibodies are monoclonal or polyclonal in origin. MART-1 protein or peptides used to generate  
30 the antibodies may be from natural or recombinant sources or generated by chemical synthesis. Natural MART-1 proteins can be isolated from mammalian biological samples. Biological samples include, but is not limited to mammalian tissues such as fresh melanoma, skin, retina,  
35 primary or continuous cultures of mammalian cells such as melanoma cultures or cultured melanocytes. The natural

° MART-1 proteins may be isolated by the same methods described above for recombinant proteins. Recombinant MART-1 proteins or peptides may be produced and purified by conventional methods. Synthetic MART-1 peptides may be custom ordered or commercially made based on the predicted amino acid sequence of the present invention (Figure 1; SEQ ID:2) or synthesized by methods known to one skilled in the art (Merrifield, R.B. (1963) J. Amer. Soc. 85:2149). Examples of MART-1 peptides include, but are not limited to, are AAGIGILTV (M9-2; SEQ. ID NO. 4), EAAGIGILTV (M10-3; SEQ. ID NO. 17) and AAGIGILTVI (M10-4; SEQ. ID NO. 18) (peptides are presented in single letter amino acid code). The most preferred peptide is AAGIGILTV (SEQ ID NO:4). If the peptide is too short to be antigenic it may be conjugated to a carrier molecule to enhance the antigenicity of the peptide. Examples of carrier molecules, include, but are not limited to, human albumin, bovine albumin and keyhole limpet hemo-cyanin ("Basic and Clinical Immunology" (1991) Stites, D.P. and Terr A.I. (eds) Appleton and Lange, Norwalk Connecticut, San Mateo, California).

Exemplary antibody molecules for use in the detection methods of the present invention are intact immunoglobulin molecules, substantially intact immunoglobulin molecules or those portions of an immunoglobulin molecule that contain the antigen binding site, including those portions of immunoglobulin molecules known in the art as F(ab), F(ab')<sub>2</sub>, and F(v). Polyclonal or monoclonal antibodies may be produced by methods known in the art. (Kohler and Milstein (1975) Nature 256, 495-497; Campbell "Monoclonal Antibody Technology, the Production and Characterization of Rodent and Human Hybridomas" in Burdon et al. (eds.) (1985) "Laboratory Techniques in Biochemistry and Molecular Biology," Volume 13, Elsevier Science Publishers, Amsterdam). The antibodies or antigen binding fragments may also be produced by genetic engineering. The technology for expression of both heavy

and light chain genes in E. coli is the subject of the PCT patent applications: publication number WO 901443, WO 901443 and WO 9014424 and in Huse et al. (1989) Science 246:1275-1281.

The antibodies of this invention may react with native or denatured MART-1 protein or peptides or analogs thereof. The specific immunoassay in which the antibodies are to be used will dictate which antibodies are desirable. Antibodies may be raised against the MART-1 protein or portions thereof or against synthetic peptides homologous to the MART-1 amino acid sequence.

In one embodiment the antibodies of this invention are used in immunoassays to detect the novel MART-1 protein in biological samples. In this method the antibodies of the present invention are contacted with a biological sample and the formation of a complex between the MART-1 antigen and antibody is detected. Immunoassays of the present invention may be radioimmunoassay, Western blot assay, immunofluorescent assay, enzyme immunoassay, chemiluminescent assay, immunohistochemical assay and the like. (In "Principles and Practice of Immunoassay" (1991) Christopher P. Price and David J. Neoman (eds), Stockton Press, New York, New York; Ausubel et al. (eds) (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New York). Standard techniques known in the art for ELISA are described in Methods in Immunodiagnosis, 2nd Edition, Rose and Bigazzi, eds., John Wiley and Sons, New York 1980 and Campbell et al., Methods of Immunology, W.A. Benjamin, Inc., 1964, both of which are incorporated herein by reference. Such assays may be direct, indirect, competitive, or noncompetitive immunoassays as described in the art (In "Principles and Practice of Immunoassay" (1991) Christopher P. Price and David J. Neoman (eds), Stockton Pres, NY, NY; Oellirich, M. 1984. J. Clin. Chem. Clin. Biochem. 22: 895-904) Biological samples appropriate for such detection assays include mammalian tissues, melanoma and melanocyte cell

° lines, skin, retina, lymph nodes, pathology specimens,  
necropsy specimens, and biopsy specimens. Proteins may be  
isolated from biological samples by conventional methods  
described in (Ausubel et al., (eds) (1987) in "Current  
Protocols in Molecular Biology" John Wiley and Sons, New  
5 York, New York).

The antibodies of this invention can therefore be  
used in immunoassays to detect MART-1 antigen or  
alteration in the level of expression of the MART-1  
antigen in biological samples isolated from mammals  
10 afflicted with a disease or disorder. Examples of  
biological samples include, but are not limited to,  
mammalian tissues, biopsy tissue samples, melanoma and  
lymph node biopsy samples, pathology and tissue samples.  
Examples of diseases that can be assessed by these  
15 immunoassays, include, but are not limited to, melanomas  
and tissues which are secondary sites for melanoma  
metastasis. By alteration in level of expression, we mean  
an increase or decrease of the MART protein or portions  
thereof relative to a control sample. Alteration is also  
20 meant to encompass substitution, deletion or addition  
mutants of the MART-1 protein. Such mutations can be  
determined by using the antibodies of this invention known  
to react with specific epitopes of the MART-1 protein and  
determining which epitopes are present relative to a  
25 control. The antibodies of this invention can therefore  
be used in an immunoassay to diagnose, assess or prognoses  
a mammal afflicted with the disease.

In a preferred embodiment, the MART-1 antibodies of  
this invention are used to assess the presence of the  
30 MART-1 antigen from a tissue biopsy of a mammal afflicted  
with melanoma using immunocytochemistry. Such assessment  
of the delineation of the MART-1 antigen in a diseased  
tissue can be used to prognose the progression of the  
disease in a mammal afflicted with the disease.  
35 Specifically the MART-1 antibodies can be used to  
characterize the radial and vertical growth phases of the

° melanoma lesion. Conventional methods for immunohistochemistry are described in (Harlow and Lane (eds) (1988) In "Antibodies A Laboratory Manual", Cold Spinning Harbor Press, Cold Spring Harbor, New York; Ausbel et al. (eds) (1987). In Current Protocols In Molecular Biology, John Wiley and Sons (New York, New York).

In another embodiment, antibodies of this invention may be used to purify the MART-1 protein or portions thereof. Immunoaffinity chromatography can be performed by conventional methods known to one skilled in the art (Ausubel et al. (eds) (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New York).

In another preferred embodiment rabbit antisera containing antibodies which specifically recognize the MART-1 protein is used to detect said protein in Western Blot Analysis. Such antisera is directed to all, or a part or parts of the MART-1 protein or synthetic peptides derived from the MART-1 protein sequence. Preferably a MART-1 synthetic peptide derived from the MART-1 predicted amino acid sequence is used (Figure 1; SEQ ID NO:2). The peptide is synthesized by standard methods on an automated peptide synthesizer and purified by high pressure liquid chromatography (HPLC) as described in Example 2. The purified peptide may be conjugated to a carrier as described in (M. Bodanszky (1984) "Principles of Peptide Synthesis," Springer Verlag, New York, New York). Using conventional methods, rabbits may be immunized with the MART-1 protein or peptide conjugated to carriers. Preferably about 0.1 to 10 (mg) of antigen in adjuvant is used, most preferably about 1 mg of antigen in adjuvant is used. The animal receives similar booster doses and antisera titer is assessed by ELISA assay. Satisfactory levels of antisera are obtained when the anti-peptide antibody titer reaches a plateau. This antibody can be used in the standard immunoassays described above.

24

°       The recombinant or natural MART-1 protein, peptides, or analogs thereof may be used as a vaccine either prophylactically or therapeutically. When provided prophylactically the vaccine is provided in advance of any evidence of melanoma. The prophylactic administration of the MART-1 vaccine should serve to prevent or attenuate melanoma in a mammal. In a preferred embodiment mammals, preferably human, at high risk for melanoma are prophylactically treated with the vaccines of this invention. Examples of such mammals include, but are not limited to, humans with a family history of melanoma, humans with a history of atypical moles, humans with a history of FAM-M syndrome or humans afflicted with melanoma previously resected and therefore at risk for reoccurrence. When provided therapeutically, the vaccine is provided to enhance the patient's own immune response to the tumor antigen present on the melanoma or metastatic melanoma. The vaccine, which acts as an immunogen, may be a cell, cell lysate from cells transfected with a recombinant expression vector, cell lysates from cells transfected with a MART-1 recombinant expression vector, or a culture supernatant containing the expressed protein. Alternatively, the immunogen is a partially or substantially purified recombinant MART-1 protein, peptide or analog thereof. The proteins or peptides may be conjugated with lipoprotein or administered in liposomal form or with adjuvant.

      While it is possible for the immunogen to be administered in a pure or substantially pure form, it is preferable to present it as a pharmaceutical composition, formulation or preparation.

      The formulations of the present invention, both for veterinary and for human use, comprise an immunogen as described above, together with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the



° other ingredients of the formulation and not deleterious to the recipient thereof. The formulations may conveniently be presented in unit dosage form and may be prepared by any method well-known in the pharmaceutical art.

5 All methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with  
10 liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired formulation.

Formulations suitable for intravenous intramuscular, subcutaneous, or intraperitoneal administration  
15 conveniently comprise sterile aqueous solutions of the active ingredient with solutions which are preferably isotonic with the blood of the recipient. Such formulations may be conveniently prepared by dissolving solid active ingredient in water containing  
20 physiologically compatible substances such as sodium chloride (e.g. 0.1-2.0M), glycine, and the like, and having a buffered pH compatible with physiological conditions to produce an aqueous solution, and rendering said solution sterile. These may be present in unit or  
25 multi-dose containers, for example, sealed ampoules or vials.

The formulations of the present invention may incorporate a stabilizer. Illustrative stabilizers are polyethylene glycol, proteins, saccharides, amino acids, inorganic acids, and organic acids which may be used either on  
30 their own or as admixtures. These stabilizers are preferably incorporated in an amount of 0.11-10,000 parts by weight per part by weight of immunogen. If two or more stabilizers are to be used, their total amount is  
35 preferably within the range specified above. These stabilizers are used in aqueous solutions at the

24

° appropriate concentration and pH. The specific osmotic pressure of such aqueous solutions is generally in the range of 0.1-3.0 osmoles, preferably in the range of 0.8-1.2. The pH of the aqueous solution is adjusted to be within the range of 5.0-9.0, preferably within the range of 6-8. In formulating the immunogen of the present invention, anti-adsorption agent may be used.

Additional pharmaceutical methods may be employed to control the duration of action. Controlled release preparations may be achieved through the use of polymer to complex or absorb the proteins or their derivatives. The controlled delivery may be exercised by selecting appropriate macromolecules (for example polyester, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and the concentration of macromolecules as well as the methods of incorporation in order to control release. Another possible method to control the duration of action by controlled-release preparations is to incorporate the MART-1 protein, peptides and analogs thereof into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating these agents into polymeric particles, it is possible to entrap these materials in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxy-methylcellulose or gelatin-microcapsules and poly(methylmethacrylate) microcapsules, respectively, or in colloidal drug delivery systems, for example, liposomes, albumin microspheres, microemulsions, nanoparticles, and nanocapsules or in macroemulsions.

When oral preparations are desired, the compositions may be combined with typical carriers, such as lactose, sucrose, starch, talc magnesium stearate, crystalline cellulose, methyl cellulose, carboxymethyl cellulose,

glycerin, sodium alginate or gum arabic among others.

The proteins of the present invention may be supplied in the form of a kit, alone, or in the form of a pharmaceutical composition as described above.

Vaccination can be conducted by conventional methods. For example, the immunogen can be used in a suitable diluent such as saline or water, or complete or incomplete adjuvants. Further, the immunogen may or may not be bound to a carrier to make the protein immunogenic. Examples of such carrier molecules include but are not limited to bovine serum albumin (BSA), keyhole limpet hemocyanin (KLH), tetanus toxoid, and the like. The immunogen also may be coupled with lipoproteins or administered in liposomal form or with adjuvants. The immunogen can be administered by any route appropriate for antibody production such as intravenous, intraperitoneal, intramuscular, subcutaneous, and the like. The immunogen may be administered once or at periodic intervals until a significant titer of anti-MART-1 immune cells or anti-MART-1 antibody is produced. The presence of anti-MART-1 immune cells may be assessed by measuring the frequency of precursor CTL (cytotoxic T-lymphocytes) against MART-1 antigen prior to and after immunization by a CTL precursor analysis assay (Coulie, P. et al., (1992) International Journal Of Cancer 50:289-297). The antibody may be detected in the serum using the immunoassay described above.

The administration of the vaccine or immunogen of the present invention may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the immunogen is provided in advance of any evidence or in advance of any symptom due to melanoma. The prophylactic administration of the immunogen serves to prevent or attenuate melanoma in a mammal. When provided therapeutically, the immunogen is provided at (or shortly after) the onset of the disease or at the onset of any symptom of the disease. The therapeutic administration of

° the immunogen serves to attenuate the disease.

A preferred embodiment is a vaccine prepared using recombinant MART-1 protein or peptide expression vectors. To provide a vaccine to an individual a genetic sequence which encodes for all or part of the MART-1 nucleic acid sequence is inserted into a expression vector, as  
5 described above, and introduced into the mammal to be immunized. Examples of vectors that may be used in the aforementioned vaccines include, but are not limited to, defective retroviral vectors, adenoviral vectors vaccinia  
10 viral vectors, fowl pox viral vectors, or other viral vectors (Mulligan, R.C., (1993) Science 260:926-932). The viral vectors carrying all or part of the MART-1 nucleic sequence can be introduced into a mammal either prior to any evidence of melanoma or to mediate regression of the  
15 disease in a mammal afflicted with melanoma. Examples of methods for administering the viral vector into the mammals include, but are not limited to, exposure of cells to the virus ex vivo, or injection of the retrovirus or a producer cell line of the virus into the affected tissue  
20 or intravenous administration of the virus. Alternatively the viral vector carrying all or part of the MART-1 nucleic acid sequence may be administered locally by direct injection into the melanoma lesion or topical application in a pharmaceutically acceptable carrier. The  
25 quantity of viral vector, carrying all or part of the MART-1 nucleic acid sequence, to be administered is based on the titer of virus particles. A preferred range of the immunogen to be administered is  $10^6$  to  $10^{11}$  virus particles per mammal, preferably a human. After immunization the  
30 efficacy of the vaccine can be assessed by production of antibodies or immune cells that recognize the antigen, as assessed by specific lytic activity or specific cytokine production or by tumor regression. One skilled in the art would know the conventional methods to assess the  
35 aforementioned parameters. If the mammal to be immunized is already afflicted with melanoma or metastatic melanoma

the vaccine can be administered in conjunction with other therapeutic treatments. Examples of other therapeutic treatments includes, but are not limited to, adoptive T cell immunotherapy, coadministration of cytokines or other therapeutic drugs for melanoma.

Alternatively all or parts thereof of a substantially or partially purified the MART-1 protein may be administered as a vaccine in a pharmaceutically acceptable carrier. Ranges of MART-1 protein to be administered are 0.001 to 100 mg per patient, preferred doses are 0.01 to 100mg per patient. In a preferred embodiment, the MART-1 peptide AAGIGILTV (SEQ. ID NO. 4) (presented in single letter code) or analogs thereof is administered therapeutically or prophylactically to a mammal in need of such treatment. Preferred doses are 0.001 mg to 100 mg, most preferred are 0.01 mg to 100 mg. The peptide may be synthetically or recombinantly produced. Immunization is repeated as necessary, until a sufficient titer of anti-immunogen antibody or immune cells has been obtained.

In yet another alternative embodiment a viral vector, such as a retroviral vector, can be introduced into mammalian cells. Examples of mammalian cells into which the retroviral vector can be introduced include, but are not limited to, primary mammalian cultures or continuous mammalian cultures, COS cells, NIH3T3, or 293 cells (ATTC #CRL 1573). The means by which the vector carrying the gene may be introduced into a cell includes, but is not limited to, microinjection, electroporation, transfection or transfection using DEAE dextran, lipofection, calcium phosphate or other procedures known to one skilled in the art (Sambrook et al. (EDS) (1989) in "Molecular Cloning. A laboratory manual", Cold Spring Harbor Press Plainview, New York). The mammalian cells expressing the MART-1 antigen can be administered to mammals and serve as a vaccine or immunogen. Examples of how the cells expressing MART-1 antigens can be administered include, but is not limited to, intravenous, intraperitoneal or intralesional.

° In a preferred embodiment, the part of the MART-1 nucleic acid sequence corresponding to the peptide AAGIGILTV (SEQ ID NO: 4) is inserted into the MART-1 expression vector and introduced into the mammalian cells.

5 The vaccine formulation of the present invention comprise an immunogen that induces an immune response directed against the melanoma associated antigens such as the melanoma associated MART-1 antigen. The vaccine formulations can be evaluated first in animal models, initially rodents, and in nonhuman primates and finally in 10 humans. The safety of the immunization procedures is determined by looking for the effect of immunization on the general health of the immunized animal (weight change, fever, appetite behavior etc.) and looking for pathological changes on autopsies. After initial testing 15 in animals, melanoma cancer patients can be tested. Conventional methods would be used to evaluate the immune response of the patient to determine the efficiency of the vaccine.

20 In yet another embodiment of this invention all, part, or parts of the MART-1 protein or MART-1 peptides may be exposed to dendritic cells cultured in vitro. The cultured dendritic cells provide a means of producing T-cell dependent antigens comprised of dendritic cell modified antigen or dendritic cells pulsed with antigen, 25 in which the antigen is processed and expressed on the antigen activated dendritic cell. The MART-1 antigen activated dendritic cells or processed dendritic cell antigens may be used as immunogens for vaccines or for the treatment of melanoma. The dendritic cells should be 30 exposed to antigen for sufficient time to allow the antigens to be internalized and presented on the dendritic cells surface. The resulting dendritic cells or the dendritic cell process antigens can then be administered to an individual in need of therapy. Such methods are 35 described in Steinman et al. (WO93/208185) and in Banchereau et al. (EPO Application 0563485A1) which are

° incorporated herein by reference.

In yet another embodiment of this invention T-cells isolated from individuals can be exposed to the MART-1 protein or portions thereof in vitro and then administered to a patient in need of such treatment in a therapeutically effective amount. Examples of where T-lymphocytes can be isolated, include but are not limited to, peripheral blood cells lymphocytes (PBL), lymph nodes, or tumor infiltrating lymphocytes (TIL). Such lymphocytes can be isolated from the individual to be treated or from a donor by methods known in the art and cultured in vitro (Kawakami, Y. et al. (1989) J. Immunol. 142: 2453-3461). Lymphocytes are cultured in media such as RPMI or RPMI 1640 or AIM V for 1-10 weeks. Viability is assessed by trypan blue dye exclusion assay. The lymphocytes are exposed to all or part of the MART-1 protein for part or all of the culture duration. In a preferred embodiment the lymphocytes are exposed to the AAGIGILTV (SEQ ID NO: 4) peptide (presented in single letter code) at a concentration of 1-10 micrograms(ug)/ml per  $10^7$  cells for all or part of the duration of lymphocyte culture. After being sensitized to the peptide the T-lymphocytes are administered to the mammal in need of such treatment. Examples of how these sensitized T-cells can be administered to the mammal include but are not limited to, intravenously, intraperitoneally or intralesionally. Parameters that may be assessed to determine the efficacy of these sensitized T-lymphocytes include, but are not limited to, production of immune cells in the mammal being treated or tumor regression. Conventional methods are used to assess these parameters. Such treatment can be given in conjunction with cytokines or gene modified cells (Rosenberg, S.A. et al. (1992) Human Gene Therapy, 3: 75-90; Rosenberg, S.A. et al. (1992) Human Gene Therapy, 3: 57-73).

In addition to use as a vaccine, the compositions can be used to prepare antibodies to MART-1 antigen, peptides

or analogs thereof. The antibodies can be used directly as anti-melanoma agents. To prepare antibodies, a host animal is immunized using the MART-1 protein, peptides or analogs thereof as the immunogen and bound to a carrier as described above for vaccines. The host serum or plasma is collected following an appropriate time interval to provide a composition comprising antibodies reactive with the immunogen. The gamma globulin fraction or the IgG antibodies can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other techniques known to those skilled in the art. The antibodies are substantially free of many of the adverse side effects which may be associated with other anti-cancer agents such as chemotherapy.

The antibody compositions can be made even more compatible with the host system by minimizing potential adverse immune system responses. This is accomplished by removing all or a portion of the Fc portion of a foreign species antibody or using an antibody of the same species as the host animal, for example, the use of antibodies from human/human hybridomas. Humanized antibodies (i.e., nonimmunogenic in a human) may be produced, for example, by replacing an immunogenic portion of an antibody with a corresponding, but nonimmunogenic portion (i.e., chimeric antibodies). Such chimeric antibodies may contain the reactive or antigen binding portion of an antibody from one species and the Fc portion of an antibody (nonimmunogenic) from a different species. Examples of chimeric antibodies, include but are not limited to, non-human mammal-human chimeras, rodent-human chimeras, murine-human and rat-human chimeras (Robinson et al., International Patent Application 184,187; Taniguchi M., European Patent Application 171,496; Morrison et al., European Patent Application 173,494; Neuberger et al., PCT Application WO 86/01533; Cabilly et al., 1987 Proc. Natl. Acad. Sci. USA 84:3439; Nishimura et al., 1987 Canc. Res. 47:999; Wood et al., 1985 Nature 314:446; Shaw et al.,



° 1988 J. Natl. Cancer Inst. 80: 15553, all incorporated herein by reference).

General reviews of "humanized" chimeric antibodies are provided by Morrison S., 1985 Science 229:1202 and by Oi et al., 1986 BioTechniques 4:214.

5        Suitable "humanized" antibodies can be alternatively produced by CDR or CEA substitution (Jones et al., 1986 Nature 321:552; Verhoeyan et al., 1988 Science 239:1534; Biedler et al. 1988 J. Immunol. 141:4053, all incorporated herein by reference).

10        The antibodies or antigen binding fragments may also be produced by genetic engineering. The technology for expression of both heavy and light chain genes in E. coli is the subject the PCT patent applications; publication number WO 901443, WO901443, and WO 9014424 and in Huse et al., 1989 Science 246:1275-1281.

15        The antibodies can also be used as a means of enhancing the immune response. The antibodies can be administered in amounts similar to those used for other therapeutic administrations of antibody. For example, 20 pooled gamma globulin is administered at a range of about 1-100mg per patient. Thus, antibodies reactive with the MART-1 antigen can be passively administered alone or in conjunction with other anti-cancer therapies to a mammal afflicted with melanoma. Examples of anti-cancer 25 therapies include, but are not limited to, chemotherapy, radiation therapy, adoptive immunotherapy therapy with TIL.

30        Alternatively, anti MART-1 antigen antibodies can be induced by administering anti-idiotypic antibodies as immunogens. Conveniently, a purified anti-MART-1 antibody preparation prepared as described above is used to induce anti-idiotypic antibody in a host animal. The composition is administered to the host animal in a suitable diluent. Following administration, usually repeated administration, 35 the host produces anti-idiotypic antibody. To eliminate an immunogenic response to the Fc region, antibodies produced

by the same species as the host animal can be used or the Fc region of the administered antibodies can be removed. Following induction of anti-idiotypic antibody in the host animal, serum or plasma is removed to provide an antibody composition. The composition can be purified as described above for anti-MART-1 antibodies, or by affinity chromatography using anti-MART-1 antibodies bound to the affinity matrix. The anti-idiotypic antibodies produced are similar in conformation to the authentic MART-1-antigen and may be used to prepare an MART-1 melanoma antigen vaccine rather than using the MART-1 protein, peptides analogs or portions thereof.

When used as a means of inducing anti-Mart-1 antibodies in an animal, the manner of injecting the antibody is the same as for vaccination purposes, namely intramuscularly, intraperitoneally, subcutaneously, interlesionally, or the like in an effective concentration in a physiologically suitable diluent with or without adjuvant. One or more booster injections may be desirable.

The MART-1 derived proteins of the invention are also intended for use in producing antiserum designed for pre- or post-disease prophylaxis. Here the MART-1 antigen, peptides or analogs thereof is formulated with a suitable adjuvant and administered by injection to human volunteers, according to known methods for producing human antisera. Antibody response to the injected proteins is monitored, during a several-week period following immunization, by periodic serum sampling to detect the presence of anti-MART-1 serum antibodies, using an immunoassay as described herein.

The antiserum from immunized individuals may be administered as a prophylactic measure for individuals who are at risk of developing melanoma. The antiserum is also useful in treating an individual afflicted with melanoma for post-disease prophylaxis.

For both in vivo use of antibodies to MART-1 antigen

and anti-idiotypic antibodies and diagnostic use, it may be preferable to use monoclonal antibodies. Monoclonal anti-MART-1 antibodies or anti-idiotypic antibodies can be produced as follows. The spleen or lymphocytes from an immunized animal are removed and immortalized or used to prepare hybridomas by methods known to those skilled in the art. (Goding, J.W. 1983. Monoclonal Antibodies: Principles and Practice, Pladermic Press, Inc., NY, NY, pp. 56-97). To produce a human-human hybridoma, a human lymphocyte donor is selected. A donor known to have a melanoma carrying the MART-1 antigen may serve as a suitable lymphocyte donor. Lymphocytes can be isolated from a peripheral blood sample or spleen cells may be used if the donor is subject to splenectomy. Epstein-Barr virus (EBV) can be used to immortalize human lymphocytes or a human fusion partner can be used to produce human-human hybridomas. Primary in vitro immunization with peptides can also be used in the generation of human monoclonal antibodies. Examples of preferred MART-1 peptides, but not limited to are, AAGIGILTV (SEQ ID NO: 4), EAAGIGILTV (SEQ ID NO: 17) and AAGIGILTVI (SEQ ID NO: 18) (peptides are presented in single letter amino acid code). Most preferably AAGIGILTV (SEQ ID NO: 4) is used as the immunogen.

Antibodies secreted by the immortalized cells are screened to determine the clones that secrete antibodies of the desired specificity. For monoclonal MART-1 antigen or peptide antibodies, the antibodies must bind to MART-1 antigen or peptide. For monoclonal anti-idiotypic antibodies, the antibodies must bind to anti-MART-1 antibodies. Cells producing antibodies of the desired specificity are selected.

The antibodies or chimeric antibodies described herein may also be coupled to toxin molecules radioisotopes and drugs by conventional methods (Vitetta et al. (1991) in "Biologic Therapy of Cancer" De Vita VT, Hellman S., Rosenberg, S.A. (eds) J.B. Lippincott Co.

Philadelphia; Larson, S.M. et al. (1991) in "Biological Therapy of Cancer" De Vita V.T., Hellman S., Rosenberg, S.A. (eds) J.B. Lippincott Co., Philadelphia). Examples of toxins to which the antibodies may be coupled to include, but are not limited to, ricin or diphtheria toxin. Examples of drugs or chemotherapeutic agents include, but are not limited to, cyclophosphamide or doxorubicin. Examples of radioisotopes, include, but are not limited to, <sup>131</sup>I. Antibodies covalently conjugated to the aforementioned agents can be used in cancer immunotherapy for treating melanoma.

Local administration to the afflicted site may be accomplished through means known in the art, including, but not limited to, topical application, injection, and implantation of a porous device containing cells recombinantly expressing the infusion, implantation of a porous device in which the MART-1 antibodies or chimeric antibodies, antibodies coupled to toxins, drugs or radiolabels or portions thereof are contained.

The above described antibodies and antigen binding fragments thereof may be supplied in kit form alone, or as a pharmaceutical composition for in vivo use. The antibodies may be used for therapeutic uses, diagnostic use in immunoassays or as an immunoaffinity agent to purify the MART-1 protein or peptides as described herein.

The present invention also provides a substantially purified and isolated nucleic acid sequence, designated c(complimentary)DNA25 (Figure 4; SEQ. ID. NO. 26) which encodes a second melanoma recognized by tumor infiltrating lymphocytes. The TIL which recognize the melanoma antigen encoded by cDNA25, are associated with in vivo tumor rejection. The TIL recognized the melanoma antigen encoded by cDNA25 in the context of HLA-A2. Comparison of the cDNA25 nucleic acid sequence (Figure 4; SEQ. ID NO. 26) with the nucleic acid sequences for genes encoding a melanocyte-melanoma specific protein gp100 shows this sequence to be similar, but distinct, from the previously

identified sequences for gp100. Previously identified sequences for gp100 include, gp100 (GenBank Accession No. M32295; also designated gp95), Pmel 17 (GenBank Accession No. M77348; Kwon et al., (1991) Proc. Natl. Acad. Sciences (USA) 88:9228-9232) and ME20 (Maresh et al. (1994) DNA and Cell Biology 13:87-95).

The cDNA25 sequence provided herein (Figure 4; SEQ. ID. NO. 26) differs from the previously reported gp100 sequence in Genbank (Genbank Accession No. M32295) by two nucleotides, from the Pmel 17 sequence (Kwon et al. (1991) Proc. Natl. Acad. Sciences (USA) 88: 9228-9232) by three nucleotides and a twenty one base pair deletion, and from the ME20 sequence (Maresh et al. (1994) DNA and Cell Biology 13:87-95) by a single nucleotide difference. At the amino acid level, the protein encoded by cDNA25 differs from the gp100 in GenBank (GenBank Accession # M32295) by one amino acid at position 162, by a two amino acid difference at positions 162 and 274, compared to Pmel 17 and did not contain 7 amino acids that existed in Pmel 17 at positions 588-594. Therefore, cDNA25 appears to encode for a different form of the gene for gp100. The differences between the cDNA25 nucleic acid sequence (Figure 4; SEQ. ID. NO. 26) and amino acid sequence (Figure 5A; SEQ. ID. NO. 27) and previously reported gp100 sequences may be due to polymorphisms, allelic variations, or to mutations within the tumor. Experiments with mouse tumors have shown that new antigens recognized by T-cells can result from point mutation in the coding region of the inactive gene (Boon, T (1992) Advances in Cancer Research 58:177-210).

This invention also provides immunogenic peptides derived from the gp100 protein sequence provided herein. (Figure 5A; SEQ. ID. NO. 27). These immunogenic peptides represent the antigenic portions of the gp100 protein (Figure 5A; SEQ. ID. NO. 27) recognized by TIL. Examples of immunogenic peptides include, but are not limited to, LLDGTATLRL (peptide G10-4; ~~SEQ ID NO: 31~~ <sup>SEQ ID NO: 31</sup>; ~~SEQ ID NO: 33~~; Figure 4,

° nucleotides 1407-1436), VLYRYGSFSV (peptide G10-5; ~~SEQ ID NO: 32~~ <sup>SEQ ID NO: 32</sup> ~~NO: 34~~; Figure 4 nucleotides 1464-1493), ALDGGNKHFL  
b (peptide G10-22; ~~SEQ ID NO: 35~~ <sup>SEQ ID NO: 33</sup> ~~SEQ ID NO: 35~~; Figure 4, nucleotides 708-  
b 737), VLKRCLLHL (peptide G9-19 ~~SEQ ID NO: 36~~ <sup>SEQ ID NO: 34</sup> ~~SEQ ID NO: 36~~; Figure 4,  
5 nucleotides 48-74), VLPSPACQLV (peptide G10-8; ~~SEQ ID NO: 37~~ <sup>SEQ ID NO: 35</sup> ~~SEQ ID NO: 37~~; Figure 4 nucleotides 1668 to 1797) SLADTNSLAV (peptide  
b G10-9; ~~SEQ ID NO: 38~~ <sup>SEQ ID NO: 36</sup> ~~SEQ ID NO: 38~~; Figure 4 nucleic acids 1746 to 1775)  
b SVSVSQLRA (peptide G10-216; ~~SEQ ID NO: 39~~ <sup>SEQ ID NO: 37</sup> ~~SEQ ID NO: 39~~; Figure 4,  
nucleotides 684 to 710) YLEPGPVTA (peptide G10-280; ~~SEQ ID NO: 40~~ <sup>SEQ ID NO: 38</sup> ~~SEQ ID NO: 40~~; Figure 4; nucleotides 876-902) or LNVSLADTN  
b 10 (peptide G10-400; ~~SEQ ID NO: 41~~ <sup>SEQ ID NO: 39</sup> ~~SEQ ID NO: 41~~; Figure 4; nucleotides  
1736-1764). This invention further includes analogs of  
these immunogenic peptides derived from the gp100 amino  
acid sequence (Figure 5A; SEQ ID NO: 27). The term analog  
15 of these immunogenic peptides. The term analog also  
includes conservative substitution or chemical derivative  
of the peptides as described above. These immunogenic  
peptides may be synthetically or recombinantly produced in  
the same manner or fashion as described above for MART-1.  
20 In another embodiment the immunogenic peptides (~~SEQ ID NO: 33 to 38~~ <sup>SEQ ID NOS: 31 to 36</sup> ~~SEQ ID NO: 33 to 38~~) derived from the gp100 amino acid  
sequence (SEQ ID NO: 27), may be used as a vaccine either  
therapeutically or prophylactically. When provided,  
prophylactically the vaccine is provided in advance of any  
25 evidence of melanoma. The prophylactic administration of  
these peptides should serve to prevent or attenuate  
melanoma in a mammal.

In a preferred embodiment, mammals preferably humans,  
at high risk for melanoma are prophylactically treated  
30 with these vaccines. Alternatively, the vaccine may be  
provided therapeutically to enhance the patients own  
immune response to the tumor antigen prescribed on the  
melanoma or metastatic melanoma. The vaccine, which acts  
as an immunogen, may be a cell, cell lysate from cells  
35 transfected with a recombinant expression vector carrying  
a nucleic acid sequences encoding gp100 immunogenic

peptide or a culture supernatant containing the expressed protein. Expression vectors into which nucleic acid sequences encoding these immunogenic peptides may be introduced are the same as those described above for MART-1. Alternatively, the immunogen is a partially or substantially purified recombinant gp100 peptide or analog thereof.

While it is possible for the immunogen to be administered in a pure or substantially pure form, it is preferable to present it as pharmaceutical compositions, formulations or preparations as described above for MART-1. Vaccination can be conducted by conventional methods previously described above for MART-1.

The gp100 immunogenic peptides and nucleic acids sequences encoding them may be used in bioassays, or to generate antibodies in the same manner or fashion as described above for MART-1.

In yet another embodiment of this invention, multivalent vaccines against one or more melanoma antigens are provided. Such multivalent vaccines may comprise all or part of the MART-1 protein peptides or gp100 peptides or combinations thereof.

Previous identification of genes encoding melanoma antigens have utilized PBL isolated from melanoma patients immunized or pretreated with antigens (Van Der Bruggen et al. (1991) Science 254: 1643-1647; Brichard et al. (1993) J. Exp. Med. 178: 489-495; Traversari, C., et al. (1992) J. Exp. Med. 176: 1453-1457). A preferred strategy is to identify genes coding for tumor antigens that are recognized by TIL from tumor-bearing patients, in the absence of immunization of said patients. Such a strategy enhances the possibility that the genes identified code for antigens involved in the natural immune response against the growing cancer. Thus, this invention also provides a method of identifying genes encoding melanoma antigens utilizing cDNA expression cloning using tumors infiltrating lymphocytes isolated (TIL) from the tumor of

patients afflicted with melanoma. The method comprises the following steps: (a) isolating tumor infiltrating lymphocytes from the tumor of a mammal afflicted with melanoma; (b) introducing a melanoma cDNA library into a mammalian cell line; (c) exposing said mammalian cells to said TIL; (d) screening for expression of an antigen encoded by said cDNA in said mammalian cells recognized by said TIL; and (e) isolating said cDNA corresponding to said antigen. The tumor infiltrating lymphocytes in step (a) may be isolated from patients afflicted with melanoma including, but not limited to, the melanoma lesion, subcutaneous tissue or visceral organs. Examples of cells that may be used to prepare the cDNA library used in step (b), include, but are not limited to, fresh or cultured melanoma cells. Preferably, the cDNA library is introduced into mammalian cells not expressing melanoma antigens. If non human mammalian cells or human cells not expressing the desired HLA haplotype for recognition by the TIL are used in step (b), such cells can be cotransfected with an HLA gene as described below. Examples of cells which can be used in step (b), include but are not limited to, tumor cell lines, such as breast cancer cell line MDA 231 (ATCC # HTB26), or COS 7 cells (ATCC #CRL 1651). Examples of MHC genes which can be used include, but are not limited to, HLA-A, HLA-B, and HLA-C genes, preferably HLA-A2 and subtypes thereof (Zemmour, J. et al. (1992) Tissue Antigens 40:221-228). The appropriate MHC gene to be used is determined by the haplotype of the tumor cells which were the source for the cDNA library. Standard methods can be used to determine the haplotype recognized by the TIL isolated (ASHI Laboratory Manual (2nd Edition 1990). Examples of how to evaluate recognition of the cells containing the cDNA clone expressing an antigen recognized by the TIL includes, but is not limited to,  $\gamma$ -interferon assays, TNF secretion (Van de Bruggen et al., (1991) Science 254:1643-1647) or lysis of cells transfected with cDNA encoding for

41



the recognized antigen. Such assays are performed by conventional methods known to one skilled in the art. Melanoma antigens can be isolated by or rescued by PCR using primer specific to flanking site of vector containing the cDNA. Examples of how to isolate the cDNA corresponding to the antigen recognized by the TIL include, but are not limited to, PCR.

Once the genes or nucleic acid sequences encoding melanoma antigens are identified, the next step is to determine the antigenic portion or epitope of the protein encoded by these genes. Therefore, in yet another embodiment of this invention, a method is provided for assessing the immunogenicity of peptides derived from the predicted amino acid sequences of the MART-1 protein (Figure 1; SEQ ID NO: 2) or a gp100 protein (Figure 5A; SEQ ID NO: 27). The method comprises the steps of: (a) preparing a plurality of peptides based on the MART-1 (Figure 1; SEQ ID NO: 2) or gp100 (Figure 5A; SEQ ID NO: 27) amino acid sequence; (b) incubating at least one of said peptides with a mammalian cell line; (c) exposing said mammalian cells incubated with said peptide to tumor infiltrating lymphocytes (TIL); and (d) screening for recognition of TIL with said cells incubated with said peptide. It is preferred that peptides of about 25 to 5 amino acids be used, more preferably 20 to 10 amino acids and most preferably 9-10 amino acids. Examples of cells that may be used in step (b) include, but are not limited to, T2 cells, (Cerundolo, V. et al. (1990) Nature, 345: 449-452) or EBV transformed B cell lines (Topalian et al. (1989) J. Immunol. 142: 3714-3725). Examples of how to assess recognition of the cells incubated with peptide include but is not limited to, <sup>51</sup>CR release cytotoxicity assay (Cerundolo, V. et al. (1990) Nature 345:449-452.) or lymphokine assays such as  $\gamma$ -IFN or TNF secretion. (Schwartzentruber, D. et al., (1991) J. of Immunology 146:3674-3681).

T cells recognize antigen complexed with MHC Class 1

° molecules. The MHC locus in all mammalian species contains numerous genes and is highly polymorphic. Different MHC molecules or haplotypes types bind different antigens. In humans the HLA complex contains the HLA-A, HLA-B and HLA-C gene loci which encode class I molecules.

5 Lymphocytes will recognize tumor antigens on the context of HLA Class 1 molecule. If the cells containing the recombinant MART-1 expression vector are to be screened by the TIL but are not human cells, such as COS cells, or do not express a desired haplotype an expression vector

10 containing an MHC Class 1 gene may also be introduced into the cells. (See Example 1) This, represents yet another alternative embodiment of the invention. Cells expressing MART-1 antigens and HLA antigens can be screened with TIL to detect the presence of tumor antigens in the context of

15 a specific MHC Class 1 restriction type. The appropriate haplotype is determined by the haplotype of the tumor from which the library is derived. Examples of MHC Class I genes that may be used include, but are not limited to, HLA-A, HLA-B and HLA-C genes. Examples of preferred MHC

20 specificities or restriction types include, but is not limited to HLA-A1, HLA-A2, such as the HLA-A2.1 subtype, or HLA-A24 (Zemmour, J. et al. (1992) Tissue Antigens 40:221-228). Most preferred is the HLA-A2.1 gene.

25 All books, articles, and patents referenced herein are incorporated by reference. The following examples illustrate various aspects of the invention and in no way intended to limit the scope thereof.

30 Cloning of a Gene Coding for a Shared  
Human Melanoma Antigen Recognized by  
Autologous T Cells Infiltrating Into Tumors

Example 1

*Generation of cytotoxic T lymphocytes (CTL) and culture of cell lines*

35 CTL were generated from excised tumor specimens by culturing a suspension of cells with 6000IU/ml of IL-2 (Cetus-Oncology Division, Chiron Corp. Emeryville, CA) for

° 30-70 days as described in Kawakami, Y., et al. (1988) J. Exp. Med. 168:2183-2191. TIL501 and TIL1235 were predominantly CD8<sup>+</sup> and were derived from the tumor specimens of patients with advanced metastatic melanoma. The CD8<sup>+</sup> T cell clone, TIL501.A42, was established by  
5 limiting dilution methods and cultured with 120 IU/ml of IL-2 plus irradiated (once a week for 4 to 6 times) autologous tumor cells.

Melanoma cell lines, 397mel, 501mel, 526mel, 537mel, 624mel, 888mel, 952mel, and Epstein-Barr virus (EBV)  
10 transformed B cell lines, 501EBVB, 836EBVB were established in our laboratory and cultured in RPMI1640 (GIBCO/Lifetechnologies, Grand Island N.Y.) medium containing 10% fetal calf serum (FCS) (Biofluids, Rockville MD). (Topalian et al., (1989) J. Immunol. 142:  
15 3714-3725) Normal cultured melanocytes, NHEM483, NHEM493, NHEM527, NHEM529, NHEM530, NHEM533, NHEM616 and NHEM680 were purchased from Clonetics, San Diego, CA, FM725, FM801, FM902 were provided by M. Herlyn, Wistar Institute, Philadelphia PA, HA002 was provided by R. Halaban, Yale  
20 university, New Haven, CT and cultured in melanocyte growth medium (MGM, Clonetics). Melanoma cell lines, C32, RPMI7951, WM115, A375, HS695T, Malme3M, colon cancer cell lines, Collo, SW480, WiDr, breast cancer cell lines, MDA231, MCF7, HS578, ZR75, neuroblastoma cell line, SK-N-  
25 SH, glioma cell lines, U138MG, HS683, H4, sarcoma cell line 143B, embryonal kidney cell line 293 transformed with adenovirus type 5 were purchased from ATCC, Rockville, MD. Renal cancer cell lines, UOK108 and UOK117 were provided M. Linehan NIH, Bethesda, MD. The small cell lung cancer  
30 cell line, H1092 was provided by J.D.Minna, University Texas Southwestern, Dallas, TX. Ewing's sarcoma cell lines, TC71, RD-ES, 6647 were provided by M. Tsokos, NIH, Bethesda, MD. The neuroblastoma cell line, SK-N-AS was provided by O.M. El Badry, NIH, Bethesda, MD. The  
35 plasmacytoma cell line, HMY-C1R and the M1 fibroblast cell line were provided by W.Biddison, NIH, Bethesda, MD.

44

Kidney epithelial cells, KAM, WLC were provided by D.J.Hazen-Martin and D.A.Sens, Medical University of South Carolina, Charleston, SC. The monkey kidney cell line, COS7, was provided by W. Leonard, NIH, Bethesda, MD.

*Cytotoxicity assay*

5       <sup>51</sup>Cr release assays were performed as described in Kawakami, Y., et al., (1988) J. Exp. Med. 168:2183-2191. Briefly, 5000 target cells labeled with <sup>51</sup>Cr were mixed with varying numbers of effector cells and incubated for 5 hours(h). Then supernatants were collected, radioactivity was measured and the percent specific lysis was calculated.

*IFN-γ release assay*

15       Fifty to one hundred thousand responder cells and 4x10<sup>4</sup>-10<sup>5</sup> stimulator cells were mixed in 300ul of AIM-V medium containing 120IU/ml IL-2 per well in a 96 flat-well microplate. After incubation for 20h, 100ul of supernatants were collected and added to an enzyme-linked immunosorbent assay (ELISA) plate (Immunoplate MaxiSorp, Nunc, Denmark) coated with anti-human IFN-γ monoclonal antibody (mAb) (Biosource, Camerillo, CA). After 20       overnight incubation at 4°C, the plates were washed 3 times and 100ul of a 1:2000 dilution of rabbit anti-human IFN-γ polyclonal antibody (Ab) (Biosource, Camerillo, CA) was added and incubated at 37°C for 2 h. Plates were 25       washed 3 times, and 100ul of a 1:2000 dilution of alkaline phosphatase labelled goat anti-rabbit IgG polyclonal antibody (Ab) (Boehringer Mannheim, Indianapolis, IN) was added. After a 1 h incubation at 37° C, 100ul of 4mg/ml p-Nitrophenyl phosphate (Sigma, St 30       Louis, MO) was added, incubated for 10-20 min at room temperature in the dark, and 25ul of 1N NaOH was added to stop the reaction. Optical density was measured at 405nm wave length and the concentration of IFN-γ was calculated in comparison to recombinant IFN-γ standards (Biogen, 35       Cambridge, MA) measured in the same assay.

*cDNA expression cloning*

A cDNA library was constructed from poly A RNA from the HLA-A2<sup>+</sup> melanoma cell line, 501mel as described in (Miki, T., et al., (1989) Gene; 83:137-146 Miki et al. (1991) Proc. Nat. Acad. Sci. USA 88:5167-5171). Briefly, first strand cDNA was synthesized with a linker primer GGACAGGCCGAGGCGGCC(T)<sub>40</sub><sup>^</sup> followed by second strand cDNA synthesis. After treatment with T4 DNA ligase, an SfiI adaptor consisting of two oligonucleotides, ~~(SEQ ID NO: 41)~~ ~~(SEQ ID NO: 42)~~ CCAATCGCGACC<sup>^</sup> and GGTCGCGATTGGTAA<sup>^</sup> was ligated to the end of the cDNA. The cDNA was digested with SfiI and the digested fragment was isolated by passing through a spun column. The cDNA was then mixed with bacteriophage  $\lambda$ pCEV27 (Miki, T. et al., (1991) Proc. Natl. Acad. Science (USA) 88: 5167-5771) vector arms prepared by SfiI digestion and in vitro packaging was performed.

To screen for melanoma antigens, 10ug of the amplified cDNA library containing approximately 10<sup>7</sup> clones was transfected into the HLA-A2<sup>+</sup> antigen non-expressing cell lines, MDA231 clone 7 and A375 clone 1-4 using a modified calcium-phosphate method (Mammalian Transfection Kit, Stratagene). After G418 (BRL, Gaithersburg, MD) selection, individual colonies were isolated and cultured in 96 well microplates and replica plates were made. A mixture of 5x10<sup>4</sup> TIL1200 and 5x10<sup>4</sup> TIL1235 was added to the wells of the microplates containing the growing transfectants that were near confluence and incubated for 20 h. Supernatants were collected and IFN- $\gamma$  was measured by ELISA.

Polymerase chain reaction(PCR) was performed to rescue the transfected genes from the genomic DNA of positive transfectants using SP6 and T7 primers which flank the inserted genes. The amplified products were cloned in the pCRII vector (Invitrogen, San Diego, CA). For cDNA clones 22 and 23, a Hind III and XhoI fragment containing the full length cDNA was subcloned into the expression vector pcDNA3 (Invitrogen, San Diego, Ca).

To test whether the cloned cDNAs encode tumor antigens, the pcDNA3 containing the cloned genes were transiently transfected into the COS7 cell line by the DEAE dextran method (Seed, B. and Aruffo, A. (1987) Proc. Natl. Acad. Sci. (USA) 84:3365-3369). Briefly,  $3 \times 10^5$

5 cells per well in 6 well plates were incubated at  $37^\circ \text{C}$  for 4 hours (h) in 0.75 ml DMEM containing 100ug of DEAE dextran (Sigma), 0.1mM chloroquine and 1ug of the pcDNA3 containing the cloned genes and/or the pcDNA-HLA-A2.1 (Zemmour, J. et al. (1992) Tissue Antigens 40: 221-228).

10 After medium was removed, 10% DMSO solution in HBSS buffer was added and incubated for 2 min. The cells were washed once with PBS and incubated in 7.5% FCS DMEM for 2 days. The 293 cell line was transiently transfected using lipofectamine (BRL, Gaithersburg, MD) according to the  
15 manufacturer's recommendation. After incubation, the ability of the transfected COS7 or 293 cells to mediate IFN- $\gamma$  release from TIL was assessed. The expression of the HLA-A2 gene was tested by flow cytometry. Stable  
20 transfectants were made by the calcium-phosphate method and individual colonies and pooled transfectants were tested for reactivity to TIL by cytotoxicity and IFN- $\gamma$  release assays.

DNA sequencing of the cloned genes was performed by dideoxy chain termination method with dGTP and 7-deaza-dGTP. DNA and protein sequences were analyzed by the GCG  
25 program with GeneBank, and EMBL Data Library nucleotide databases and SWISS-PROT, PIR, GenPept, Brookhaven Protein Data Bank protein databases.

#### 30 *Northern blot analysis*

Total RNA was isolated by the guanidine-isothiocyanate-cesium chloride centrifugation method. (Chirgwin, J.M. et al. (1979) Biochemistry 18: 5294). Total RNA from normal tissue was purchased from Clontech,  
35 (Palo Alto, CA). Ten to twenty micrograms of total RNA was subjected to electrophoresis in a 1% agarose

formaldehyde gel and transferred to a nylon membrane (Duralon-UV membranes, Stratagene, La Jolla, CA) The Sal I digested fragment containing the full length cDNA from clone 22 and the  $\beta$ -actin cDNA (Clontech) were labeled by random priming and used as a probe. Hybridization with the probe was performed according to the QuikHyb protocol (Stratagene) at 68 C for 2-16 h. Membranes were washed two times with 2XSSC/0.1%SDS at 60 C for 15 minutes (min) and once with 0.1XSSC at 60 C at for 30 min and then autoradiography was performed.

10     *Characterization of cultured TIL from melanoma patients*

Multiple TIL lines were established from HLA-A2<sup>+</sup> melanoma patients and tested for lysis of melanoma cell lines from HLA-A2<sup>+</sup> and HLA-A2<sup>-</sup> patients. HLA typing of patients is performed by conventional HLA typing techniques. HLA-A2 was selected because it is the most frequently expressed class I MHC antigen (about 50% of individuals) and has been shown to be a dominant restriction element for the recognition of melanoma antigens (Crowley, N.J., et al. (1991), J. Immunol. 146:1692-1694). TIL501, TIL1235 and TIL1200 exhibited specific recognition of shared melanoma antigens in an HLA-A2 restricted fashion. TIL501.A42 was a T cell clone established from TIL501 by limiting dilution. These TIL caused lysis or released cytokines including IFN- $\gamma$ , TNF $\alpha$  and GM-CSF when cocultured with a variety of HLA-A2<sup>+</sup> melanoma or melanocyte cell lines but not HLA-A2<sup>-</sup> melanoma lines or HLA-A2<sup>+</sup> non-melanoma cell lines including the breast cancer cell line, MDA 231. Two representative experiments are shown in Table 1. Thus, these CTL seemed to recognize a non-mutated peptide derived from a melanocyte lineage specific antigen.

*Cloning of cDNA coding for melanoma  
antigens recognized by T cells*

35     A cDNA library from the HLA-A2<sup>+</sup> 501mel melanoma cell line was transfected into two highly transfectable HLA-

A B  
° A2.1<sup>+</sup> cancer cell lines, MDA231 and A375. These cell lines were not lysed by melanoma specific TIL but were lysed by HLA-A2 restricted influenza M1 specific CTL after incubation with the M1<sub>58-66</sub> peptide (GILGFVFTL; single letter code) derived the influenza matrix protein or infection with a recombinant vaccinia virus containing the M1 gene (data not shown). Thus, these cell lines exhibited normal antigen processing and presenting ability but were not lysed by these melanoma-specific TIL because of the lack of expression of the relevant melanoma antigens. After selection with G418, approximately 6700 transfected clones from each cell line were isolated and grown in microplates. Using the IFN- $\gamma$  release assay, 21 MDA231 and 27 A375 positive clones were isolated and rescreened. Of these clones, eight MDA231 and seven A375 clones were positive in a second screening assay.

In order to rescue the integrated genes, PCR using genomic DNA from these positive transfectants was performed with SP6 and T7 primers flanking the insert genes. Eight genes that were amplified from the seven transfectants which showed 1 to 2 sharp bands, including a 1.6Kb band from MDA-22 and MDA-23 transfectants, were subcloned into the pCRII cloning vector, and then further cloned into the pcDNA3 eukaryotic expression vector. The 1.6Kb band detected by Northern blot analysis with the cDNA 22 probe suggested that this fragment was a full length cDNA.

Transient transfection of the expression vector pcDNA3 containing the cDNA from clones 22 or 23 into either COS7 or 293 cells along with the HLA-A2.1 gene conferred reactivity to TIL1235 and TIL501.A42 as demonstrated by the specific release of IFN- $\gamma$  (Table 2, Experiments 1 and 2). Stable transfection of these cDNA fragments into MDA 231 or A375mel cell lines also conferred reactivity to TIL1235 and TIL501.42 (Table 2, Experiment 3). TIL501.A42 could lyse MDA231 stably transfected with cDNA 22 (data not shown). These results

49



- ° indicated that these cDNAs encode a melanoma antigen recognized by HLA-A2 restricted TIL from melanoma patients. Transfection of another clone, MDA-25 stimulated the release of interferon- $\gamma$  only from TIL 1200.

5

10

15

20

25

30

35

35 30 25 20 15 10 5 0

Table 1. Specificity of TIL 501.A42 and TIL 1235 for Melanomas and Melanocytes

A. Lytic Specificity*		B. Cytokine Release Specificity**					
Target	HLA-A2	TIL501.A42	TIL1235	LAK <sup>+</sup>	Stimulator <sup>+</sup> Cells	HLA-A2	TIL501.A42 TIL1235 TIL586 <sup>++</sup>
501mel	+	54	51	78	501mel	+	647 219 <50
526mel	+	25	33	74	586mel	-	<50 850 1034
624mel	+	23	27	75	NHEM493	+	1835 <50 <50
952mel	+	10	11	75	NHEM527	+	1638 <50 <50
Malme3M	+	36	41	70	NHEM530	+	1224 2532 <50
C32	+	17	23	82	NHEM533	+	251 <50 <50
RPMI7951	+	1	6	67	NHEM616	+	635 423 <50
WM115	+	-2	3	68	FM725	+	5975 1471 <50
HS695T	+	1	2	87	FM801	+	1375 893 62
397mel	-	-1	0	70	NHEM483	-	<50 <50 <50
MDA231	+	0	3	94	NHEM680	-	<50 548 <50
					HA002	-	<50

TIL 501.A42 and TIL1235 lysed most HLA-A2 melanoma cell lines and secreted IFN-γ when cultured with HLA-A2 melanomas and melanocytes.

\*51Cr release assay was performed at E:T = 20:1 for TIL501.A42, at 40:1 for TIL1235. All targets were melanoma cell lines except for MDA231 which was a breast cancer cell line.

\*\*IFN-γ in the supernatant was measured after TIL and stimulator cells were coincubated for 20 hr. 501mel and 586mel are melanoma cell lines. All others were normal melanocyte cell lines.

\*LAK: Lymphokine activated killer cells.

\*\*TIL586 is class I MHC restricted melanoma specific TIL, not restricted by HLA-A2.

Characterization of this cDNA revealed it to be similar, but distinct, from a previously described melanoma antigen gp100 recognized by monoclonal antibody HMB45. This clone is described in more detail in Example 3.

The cDNA sequence of clones 22 and 23 were identical except at a single base that was believed to be a change introduced by PCR. Two other independently amplified fragments were also sequenced to clarify this region and the consensus sequence is shown in Figure 1. The longest open reading frame in this gene consists of 354 bases corresponding to a 118 amino acid protein of 13kd. This sequence did not show significant similarity to any complete nucleotide or protein sequences in established databases. Amino acids 27-47 consist of a hydrophobic region that may contain the HLA-A2 binding peptides (Falk, K., et al. (1991), Nature 351:290-296; Hunt, D.F., et al. (1992), Science 255:1261-1263; Ruppert, J., et al. (1993), Cell 74:929-937; Nijman, H.W., et al. (1993), Eur. J. Immunol. 23:1215-1219). The antigen encoded by the cDNA 22 and 23, was designated the MART-1 antigen (Melanoma Antigen Recognized by T cells-1). Of the ten HLA-A2 restricted TIL lines generated nine recognized MART-1, and four recognized a form of gp100 isolated and described herein (see Example 3) and none appeared to recognize MAGE-1 (Zakut, R., et al. (1993), Cancer Res 53:5-8.; data not shown).

30

35

52

T530x

Table 2. Interferon Gamma Secretion by TIL501.A42 and TIL 1235 When Cultured with HLA-A2<sup>+</sup> Cell Lines Transfected with the Gene 22 or 23

Transfected Gene HLA-A2 TIL501.A42 TIL1235

Stimulator Cell	Transfected Gene	pg/ml
Exp. 1		
501mel	none	1076
397mel	none	<50
COS7	none	<50
COS7	HLA-A2.1	<50
COS7	22	<50
COS7	HLA-A2.1+22	1049
Exp. 2		
501mel	none	1051
397mel	none	<50
293	HLA-A2.1	<50
293	22	<50
293	HLA-A2.1+22	255
Exp. 3		
501mel	none	1056
397mel	none	<50
MDA231	none	<50
MDA231	23	725
A375	none	<50
A375	23	131

IFN-γ in the supernatant was measured after TIL were coinocubated for 20 hr with COS7 or 293 cell lines transiently transfected with the pCDNA3 containing the HLA-A2.1 and/or cDNA 22 by the DEAE-dextran method (Exp. 1 & 2), or with the A375 or MDA231 cell lines stably transfected with cDNA 23 (Exp. 3).

IFN-γ was secreted only when TIL were incubated with HLA-A2<sup>+</sup> cell lines transfected with the cDNA 22 or 23.

\*IFN-γ secreted by TIL alone without stimulator (<50 pg/ml) was subtracted.

+Not done

53

7540X

35

30

25

20

15

10

5

Table 3. Expression of the MART-1 Gene in a Variety of Tissue and Cell Lines

<u>Melanoma</u>		<u>Normal fresh tissue</u>	<u>Colon cancer</u>
<u>HLA-A2+</u>		Brain	Collo
501mel	+	Retina	SW480
526mel	+	Adrenal gland	WiDr
624mel	+	Lung	
Malm3M	+	Liver	<u>Breast cancer</u>
952mel	+	Kidney	MDA231
697mel	+	Spleen	MCF7
C32	+	Thymus	HS578
RPMI7951	+	Testis	ZR75
WM115	-	Fetal liver	
A375	-		<u>Neuroblastoma</u>
<u>HLA-A2-</u>		<u>Non-melanoma cell lines</u>	SK-N-AS
397mel	+	T-cell	SK-N-SH
888mel	+	TIL	<u>Ewing sarcoma</u>
537mel	+	TILB	TC75
586mel	+	B-cell	RD-ES
		Daudi	6647
		HMY-C1R	
		501EBVB	<u>Sarcoma</u>
		836EBVB	143B
<u>Melanocyte</u>			
NHEM483	+		
NHEM493	+		
NHEM529	+		
NHEM530	+		
FM902	+		
FM906	+		
HA002	+		
		<u>Kidney epithelial cell</u>	
		KAM	<u>Renal cancer</u>
		WLC	UOK108
			UOK117
			<u>Small cell lung cancer</u>
			H1092

Northern blot analysis with 10-20 ug of total RNA was probed with the full length cDNA of the gene 22. The RNA from most melanomas, all melanocyte cell lines tested and retina were positive.

*Expression of MART-1*

Northern blot analysis of a variety of cell lines including melanoma, melanocyte and nonmelanoma cancer cell lines and normal tissues was performed to evaluate the expression of the gene coding for MART-1 (Table 3). Seven of ten HLA-A2<sup>+</sup> melanoma cell lines, all four HLA-A2<sup>-</sup> melanoma cell lines, and all seven melanocyte cell lines tested were positive for MART-1 RNA expression. In this Northern analysis, all HLA-A2<sup>+</sup> melanoma cell lines recently established in our laboratory expressed MART-1 RNA. There was a perfect correlation between MART-1 expression and lysis by TIL501.A42 in the 10 HLA-A2<sup>+</sup> melanoma lines shown in Table 3. TIL 501.A42 which recognized the MART-1 Ag lysed 13 of 17 (76%) HLA-A2<sup>+</sup> melanoma cell lines tested (data not shown). Of ten normal human tissues examined for mRNA expression by Northern blot analysis only retina was positive. No positivity was seen in any cell lines from T cells, B cells, kidney epithelial cells or fibroblasts or in 19 nonmelanoma tumors. It thus appears that MART-1 is a previously undescribed antigen expressed on melanocyte lineage cells from skin and retina that is also expressed on melanoma cells.

Studies using a panel of T cell clones and immunoselected melanoma clones (Knuth, A., et al. (1989), Proc. Natl. Acad. Sci. (USA) 86:2804-2808; Wolfel, T., et al. (1987), J. Exp. Med. 170:797-810), as well as studies analyzing HPLC fractionated peptides from melanoma cells (Slingluff, C.L., et al. (1993), J. Immunol. 150:2955-2963; Storkus, W.J., et al. (1993), J. Immunol. 151:3719-3727) suggest that multiple antigenic peptides that can provoke an immune response exist on melanomas. By cDNA cloning, two genes encoding melanoma antigens have been identified; MART-1 (Figure 1; SEQ ID NO. 1) and a gp100 gene (see Example 3; Figure 4; SEQ ID NO. 26). MART-1 and the form of gp100 identified herein (Figures 4 & 5A; SEQ. ID NOS. 26 and 27), both are recognized by HLA-A2.1

55

° restricted TIL. The MART-1 antigen is a 118 amino acid protein of approximately 13 kd. Neither the gene nor the amino acid sequence for MART-1 have been previously described.

5 MART-1 RNA was expressed in 11 of 14 HLA-A2.1 positive or negative melanoma lines, and 7 of 7 melanocyte lines. With the exception of retinal tissue no MART-1 expression was found on any normal tissue tested, T-cell lines, B-cell lines, kidney epithelial lines, a fibroblast line or 19 tumor cell lines from cancers of the colon,  
10 breast, brain, kidney, lung or bone.

Another melanoma antigen, MAGE-1, has been described that is recognized by T cells derived from peripheral blood lymphocytes following repetitive in vivo or in vitro immunization (Van Der Bruggen, et al. (1991), Science  
15 254:1643-1647).

The identification of genes associated with melanoma tumor antigens opens new possibilities for active specific immunization approaches to the immunotherapy of patients with cancer based on the introduction of these genes into  
20 viral or bacterial vector systems. The possibility exists that immune reactions induced against melanocyte-melanoma lineage antigens such as MART-1 may be generated against normal cells. Vitiligo, probably resulting from anti-melanocyte immune reactions, has been reported to be  
25 associated with a favorable prognosis in patients with melanoma (Nordlund, J.J., et al. (1983), J. Am. Acad. Dermatol. 9:689-695); Bystryrn, J-C, et al. (1987), Arch. Dermatol. 123:1053-1055), and has also been reported in patients responding to chemoimmunotherapy (Richards, J.M.,  
30 et al. (1992), J. Clin. Oncol. 10:1338-1343). TIL with anti melanocyte-melanoma reactivities have been administered to patients with advanced melanoma (Rosenberg, S.A., et al. (1988), N Engl J Med 319:1676-1680; Rosenberg S.A., J. Clin. Oncol. 10:180-199) and  
35 although sporadic vitiligo has been seen in these patients, no adverse ophthalmologic effects related to the

56

possible expression of these melanocyte antigens on retinal cells has been observed.

Because HLA-A2 is present in about 50% of individuals and the HLA-A2 restricted MART-1 antigen also appears to be widely expressed on melanomas, immunization with the MART-1 antigen may be particularly useful for the development of active immunotherapies.

#### Example 2

##### Characterization Of Immunogenic Epitopes of MART-1

*Generation of melanoma specific CTL lines and a clone from TIL*

Melanoma specific CTL lines were generated by culturing a single cell suspension made from metastatic melanoma with 6000U/ml of IL2 (Cetus-Oncology Division, Chiron Corp. Emeryville, CA) as previously reported (Kawakami, Y. et al., (1988) J. Exp. Med. 168:2183) A T-cell clone, A42 was established by limiting dilution methods from patient, 501.

##### *Assessment of antigen recognition by CTL*

<sup>51</sup>Cr release cytotoxicity assays and cytokine release assays using ELISA to measure IFN- $\gamma$ , GM-CSF and TNF- $\alpha$  were performed to analyze the reactivity of TIL as described in Kawakami, Y. et al. (1988), J. Exp. Med. 168:218 (see Example 1). Melanoma cell lines were established in the laboratory. For analysis of the recognition of known antigens by TIL, the COS7 cell line transfected with cDNAs encoding either, MART-1, gp100, or tyrosinase related protein (gp75) (Cohen, T. et al., (1990) Nucleic Acids Research 18:2807) along with HLA-A2.1 cDNA were incubated with TIL for 20h and the amount of IFN- $\gamma$  secreted into the supernatant was measured by ELISA as described in Example 1. The cDNA encoding MART-1 (see Example 1) or gp100 (see Example 3) in plasmid pcDNA3 (Invitrogen, San Diego, CA) was cloned from a 501mel melanoma cDNA library by



screening with TIL1235 or TIL1200, respectively (see Example 1). The cDNA encoding tyrosinase related protein(gp75) in pCEV27 plasmid was isolated from 501mel melanoma cDNA library using a probe generated by PCR based on the reported sequence of gp75 (Cohen et al. (1980) Nucleic Acids Research 18:2807).

#### *Peptide synthesis and identification of antigenic peptides*

Peptides were synthesized by a solid phase method using a Gilson AMS 422 multiple peptide synthesizer. The peptides were purified by HPLC on a Vydac C-4 column with 0.05% TFA/water-acetonitrile. To identify the antigenic peptides, TIL lysis of T2 cell lines preincubated for 2h with each peptide was measured using a <sup>51</sup>Cr release cytotoxicity assay.

#### *HLA-A2 Restricted Melanoma Specific TIL*

HLA-A2 restricted melanoma specific CTL lines and a clone, A42, were established from lymphocytes infiltrating into tumors of 10 melanoma patients. These TIL recognized autologous and most allogeneic fresh or cultured melanoma cells expressing HLA-A2, but did not recognize HLA-A2<sup>-</sup> melanomas or HLA-A2<sup>+</sup> non-melanoma cell lines (Kawakami et al. (1992) J. Immunol 148:638). They also recognized HLA-A2<sup>+</sup> normal cultured melanocytes derived from neonatal skin (see Example 1 and Kawakami, Y. et al. (1993), J. Immunotherapy 14:88). Thus, these TIL recognized non-mutated self-peptides derived from proteins expressed in melanoma and melanocytes in association with HLA-A2.

#### *Recognition of Additional melanoma proteins by TIL*

To evaluate the frequency of recognition of 4 isolated melanoma proteins including MART-1, a form of gp100 (Figure 5A; SEQ ID NO: 26, see Example 3), and tyrosinase related protein(gp75), the reactivity of TIL to COS7 was tested on cell lines transfected with cDNAs

58

° encoding these 3 proteins with or without the cDNA  
encoding HLA-A2.1. One of several experiments with 9 TIL  
is shown in Table 4. Eight of the nine HLA-A2 restricted  
melanoma specific TIL secreted IFN- $\gamma$  when incubated with  
COS7 cotransfected with MART-1 and HLA-A2.1. Only TIL1200  
5 which is a relatively oligoclonal CTL line (Shilyansky, J.  
et al., (1994) Proc. Natl. Acad. Sci. (USA) 91:2829) did  
not respond to this COS transfectant. Four TIL (620, 660,  
1143, 1200) recognized gp100 when transfected along with  
HLA-A2.1. TIL1200 secreted large amounts of IFN- $\gamma$   
10 compared to TIL620, 660, and 1143, suggesting that only a  
small subset of T-cells in these latter 3 TIL lines  
recognized gp100. None of these TIL recognized gp75 using  
this assay. Thus, MART-1 is a common melanoma antigen  
recognized by most HLA-A2 restricted TIL derived from  
15 melanoma patients.

*Identification of MART-1 epitopes for TIL.*

To identify the MART-1 epitopes for these TIL, 23  
peptides were selected based on the known peptide binding  
20 motifs to HLA-A2.1 (Falk, K. et al., (1993) Nature,  
351:290; Hunt, D. F. et al. (1992), Science, 255:1261;  
Ruppert, J. et al., (1993) Cell 74:929), synthesized (>90%  
purity) and screened by testing lysis of the HLA-A2.1<sup>+</sup> T2  
cell line by TIL after incubation of the T2 line with each  
25 peptide (Table 5). The T2 cells (Cerundolo, V. et al.;

30

35

T600X

35 30 25 20 15 10 5 0

Table 4. Recognition of melanoma antigens by HLA-A2 restricted melanoma specific TIL

Stimulator Cell line	Transfected CDNA	HLA-A2	TIL 501	TIL 620	TIL 660	TIL 1074	TIL 1088	TIL 1128	TIL 1143	TIL 1200	TIL 1235
501mel	none	+	93	720	530	670	491	272	354	736	750
397mel	none	-	0	0	0	0	0	0	0	0	0
COS7	none	-	0	0	0	0	0	0	14	0	0
COS7	HLA-A2.1	+	0	0	0	0	0	0	0	0	0
COS7	MART-1	-	0	0	0	0	0	0	12	0	0
COS7	gp100	-	0	0	0	0	0	0	14	0	0
COS7	gp75	-	0	0	0	0	0	0	0	0	0
COS7	HLA-A2.1+MART-1	+	270	196	131	625	328	52	184	0	743
COS7	HLA-A2.1+gp100	+	0	89	17	0	0	0	41	391	8
COS7	HLA-A2.1+gp75	+	0	0	0	11	0	0	7	0	0

IFN- $\gamma$  secretion(pg/ml)

IFN- $\gamma$  was measured in the supernatants after HLA-A2 restricted melanoma specific TIL were coinocubated with COS7 cells cotransfected with cDNAs encoding proteins expressed in melanoma with or without HLA-A2.1 cDNA. All TIL except TIL1200 secreted IFN- $\gamma$  when cultured with COS7 cotransfected with cDNAs encoding MART-1 and HLA-A2.1. TIL620, 660, 1143 and 1200 secreted IFN- $\gamma$  when cultured with COS7 cotransfected with the cDNAs encoding gp100 and HLA-A2.1.

° (1990) Nature 345: 449-452) cell line was lysed well by all 4 HLA-A2 restricted melanoma specific TIL tested when preincubated with either peptides M9-2, M10-3, or M10-4. Both 10 amino acid peptides, M10-3 and M10-4 contain the M9-2 sequence, with M10-3 having an additional glutamic acid at its N-terminus and M10-4 having an extra isoleucine at its C-terminal end. These peptides are located in a hydrophobic putative transmembrane domain in MART-1. The same lysis was observed when other HLA-A2<sup>+</sup> cells incubated with these peptides were used as targets including the K4B (provided by Dr. William Biddson, NIH; Storkus, W et al. (1993) J. of Immunology 151:3719-3727) and 501EBVB Epstein-Barr virus transformed B cells (Topalian et al. (1989) J. Immunol. 142: 3714-3725) or HMY-C1R B cells (Dr. William Biddson; NIH; Storkus, W. et al., (1993) J. of Immunol. 151:3719-3727) transfected with the HLA-A2.1 gene (data not shown).

The peptides, M9-1, M9-2, M9-3, M10-2, M10-3, M10-4 and M10-5 were further purified and titrated in order to evaluate their relative ability to sensitize T2 cells to lysis by MART-1 reactive TIL1235 or T cell clone A42 (Figure 2). The purified peptides M9-2, M10-3 and M10-4 were required in minimum concentrations of 1ng/ml, 100ng/ml and 1000ng/ml, respectively. The purified M10-4 was not recognized by TIL clone A42 even at 10ug/ml as shown in Figure 2. M9-1, M9-3, M10-2, and M10-5 peptides were not recognized by either A42 or TIL1235.

30

35

Table 5. Lysis of T2 cells preincubated with synthetic MART-1 peptides

Target	Peptide	A42	TIL 1235	TIL 660	TIL 1074
% specific lysis					
501mel	none	47	30	31	41
397mel	none	1	0	1	2
T2	none	-2	-3	-1	1
T2	M9-1 TTAEAAAGI	-10	-5	-5	-4
T2	M9-2 <u>AAGIGILTV</u>	64	80	40	56
T2	M9-3 GIGILTVIL	18	20	0	10
T2	M9-4 GILTVILGV	1	-1	-3	2
T2	M9-5 ILTVILGVL	-2	-1	-5	-1
T2	M9-6 LTVILGVLL	1	0	1	0
T2	M9-7 TVILGVLLI	-2	-3	-2	1
T2	M9-8 VILGVLLLI	1	5	-2	-2
T2	M9-9 ALMDKSLHV	-1	-4	-8	0
T2	M9-10 SLHVGQTQCA	-1	1	-8	4
T2	M9-11 PVVPNAPPA	-2	0	4	-1
T2	M9-12 NAPPAYEKL	1	-2	0	6
T2	M10-1 YTTAEAAAGI	-4	-2	-3	3
T2	M10-2 TAEAAAGIGI	7	11	12	15
T2	M10-3 <u>EAAGIGILTV</u>	55	66	31	51
T2	M10-4 <u>AAGIGILTVI</u>	34	68	29	21
T2	M10-5 GILTVILGVL	-1	2	7	10
T2	M10-6 ILTVILGVLL	1	6	6	7
T2	M10-7 LTVILGVLLI	-2	-1	-1	2
T2	M10-8 TVILGVLLLI	-6	-1	-1	11
T2	M10-9 RALMDKSLHV	3	5	8	11
T2	M10-10 SLHVGQTQCAL	-2	-8	2	9
T2	M10-11 SLQEKNCPEV	3	2	2	9

Twenty-three peptides (12 9-mers and 11 10-mers) (>90% purity) were synthesized and the lysability by TIL clone A42, TIL lines TIL1235, TIL660, and TIL1074 derived from different patients was tested against HLA-A2<sup>+</sup> T2 cells preincubated with each peptide (10ug/ml) in a 4h-<sup>51</sup>Cr release cytotoxicity assay at E:T ratio of 20:1 for A42 and 40:1 for other TIL lines. T2 cells were lysed well when incubated with M9-2, M10-3 and M10-4. M10-3 and M10-4 contain the entire M9-2 sequence (underlined).

*Recognition of MART-1 peptides  
by HLA-A2 restricted TIL  
established from different patients.*

To evaluate whether a variety of HLA-A2 restricted  
5 MART-1 specific TIL recognized the same or different  
epitopes in the MART-1 antigen, lysis of T2 cells  
(Cerundolo V., et al. (1990) Nature 345: 449-452)  
preincubated with each peptide was tested with TIL derived  
from 10 melanoma patients. A representative experiment  
10 with 10 TIL is shown in Table 6. M9-2 and M10-3 were  
recognized by 9 of 10 TIL (only TIL1200 were negative) as  
well as the A42 clone with the same pattern of lysis as  
COS7 cells cotransfected with cDNAs encoding MART-1 and  
HLA-A2.1. Only TIL620 and TIL1088 demonstrated low level  
15 of non-specific lysis of T2 cells without peptides or  
after the addition of irrelevant peptides, but showed  
significant increase of lysis of T2 cells preincubated  
with M9-2, M10-3, and M10-5 peptides. The recognition of  
M10-4 differed among the TIL, but was similar to the  
20 different reactivity to M10-4 by the T-cell clone A42 or  
the T-cell line TIL1235 (Figure 2). Higher concentrations  
(1ug/ml) of M10-4 were required for lysis than were  
required for M9-2 or M10-3. These 10 TIL and clone A42  
also secreted cytokines including IFN- $\gamma$ , GM-CSF and TNF- $\alpha$   
25 when incubated with T2 cells preincubated with M9-2 or  
M10-3 (data not shown). Therefore, M9-2 or M10-3 are  
common epitopes recognized by a majority of HLA-A2  
restricted melanoma specific TIL.

30

35

Table 6. Recognition of MART-1 peptides by HLA-A2 restricted melanoma specific TIL

Target Peptide (ug/ml)	TIL 501	TIL 620	TIL 660	TIL 1074	TIL 1088	TIL 1128	TIL 1143	TIL 1200	TIL 1235	TIL 1363	clone A42
---------------------------	------------	------------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------

## % specific lysis

501mel	none	42	49	35	32	31	19	24	41	32	43	41
397mel	none	3	16	6	1	1	4	3	3	3	4	1
T2	none	0	7	-3	-6	7	-6	-7	-6	-7	-7	-6
T2	M9-1 (1)	4	15	-4	1	31	1	-5	-1	1	4	3
T2	M9-2 (1)	86	75	73	79	98	30	36	2	92	82	91
T2	M9-2 (0.001)	52	49	23	32	81	9	6	1	10	41	63
T2	M9-3 (1)	5	25	0	1	19	0	1	-2	0	-2	-4
T2	M10-2 (1)	10	22	5	8	21	8	3	7	7	7	6
T2	M10-3 (1)	84	68	68	73	79	24	27	1	42	67	62
T2	M10-3 (0.001)	91	50	33	25	86	13	14	0	14	39	1
T2	M10-4 (1)	83	47	16	35	80	6	3	1	14	53	0
T2	M10-4 (0.001)	0	11	3	0	14	4	-1	-1	2	-3	-3
T2	M10-5 (1)	4	14	1	4	13	2	3	0	3	0	2

Lysability by TIL clone A42 and TIL lines derived from 10 patients of T2 cells preincubated with the purified peptides M9-1, M9-2, M9-3, M10-2, M10-3, M10-4, and M10-5 was tested in a 4h-<sup>51</sup>Cr release assay at an E:T ratio of 20:1 for A42 and 40:1 for other TIL lines. Nine of ten TIL lysed T2 cells incubated with peptides M9-2 or M10-3. Seven of ten TIL lysed T2 incubated with peptide M10-4 at a concentration of 1ug/ml.

The relative frequency of recognition of known melanoma proteins by T-cells derived from the TIL of ten melanoma patients has been examined. The common epitopes, M9-2 and M10-3 in the MART-1 antigen that were dominantly recognized by nine of these TIL have also been identified. The cDNA encoding MART-1 was isolated by cDNA expression cloning using TIL1235 in screening assays (See Example 1). MART-1 is a 118 amino acid protein containing a single transmembrane domain and is expressed in most melanoma cells as well as cultured melanocytes and retina similar to the expression pattern of the cDNA for a form of gp100 described in Example 3. The gp100 is recognized by 4 of 10 TIL.

Based on dose response analysis, peptide M9-2 most effectively sensitized T2 cells for lysis (Figure 2) suggesting that this peptide may be naturally processed and presented on tumor cells. The T-cells recognizing M9-2 may react with peptide M10-3 or M10-4 because the latter 10-mer peptides contain the 9 amino acid sequence of peptide M9-2. There is some difference in recognition of these 3 peptides by different TIL. For example, M10-4 was poorly recognized by the T-cell clone A42, but was well recognized by some TIL lines, although a higher concentration of M10-4 was necessary to observe the lysis. This may be due to the variation of TCR affinity for the M9-2 and M10-4 peptides in the context of HLA-A2, or alternatively, TIL lines may contain different T-cell clones which only recognize either M9-2 or M10-4. Peptides M10-3 and M10-4 may also be naturally processed and presented by tumor cells. The existence of multiple melanoma antigens presented by HLA-A2 has previously been suggested by analyzing the recognition of melanoma cell clones by a variety of T-cell clones (Knuth, A. et al. (1989), Proc. Natl. Acad. Sci. (USA) 86:2804, Wolfel, T. et al., 1989 J. Exp. Med. 170:797) or by analyzing HPLC peptide fractions that were isolated from HLA-A2 melanoma cells (Slingluff, C. L. Jr. et al., (1993) J. Immunol.



150:2955, Storkus; W. J. et al., (1993) J. Immunol.  
151:3719).

The observation that most HLA-A2 restricted TIL from melanoma patients recognize common MART-1 peptides but not gp75 suggests that the M9-2 or M10-3 MART-1 peptides may be more immunogenic in inducing T-cell responses in vivo than other known melanoma antigens. Some of the TIL used in this study were injected along with IL2 into autologous patients, and interestingly, all 4 TIL (620, 660, 1074, 1200) that recognize a gp100 protein (Figure 5A; SEQ ID NO: 27) effectively induced tumor regression (more than 50% reduction of tumor). All but TIL1200 also recognized MART-1.

### Example 3

#### Identification of a Second Human Melanoma Antigen Recognized by Tumor Infiltrating Lymphocytes Associated with in Vivo Tumor Rejection

##### *cDNA expression cloning*

The cDNA25 clone encoding a form of the melanoma antigen designated gp100 was cloned by techniques similar to those described in Example 1 and in Miki, T., et al. (1991) Proc. Natl. Acad. Sci. (USA) 88:5167-5171. Briefly, a breast cancer cell line, MDA231 (ATCC #HTB26), transfected with a cDNA library in  $\lambda$ pCEV27 made from the 501mel melanoma cell line was screened for antigen positivity by measuring interferon- $\gamma$  (IFN- $\gamma$ ) secretion when cocultured with TIL1200. TIL1200 was generated as described in Kawakami, Y., (1988), J. Exp. Med. 168, 2183-2191. The integrated cDNA was recovered from the genomic DNA of positive transfectants by PCR and cloned into the mammalian expression plasmid pCDNA3 (Invitrogen, San Diego, CA). The full length cDNA for cDNA25 was isolated from the 501mel  $\lambda$ pCEV27 library using the cDNA25 probe. The  $\lambda$ phage containing the full length cDNA25 was digested with XhoI, and then self-ligated with T4 DNA ligase to make the plasmid pCEV27-FL25. Alternatively, a full

length cDNA25 isolated by PCR using the specific primers designed for gp100 was cloned in pCRII(Invitrogen), and then cloned into pcDNA3(pcDNA3-FL25). To test whether this cDNA encoded a melanoma antigen it was retransfected into COS7, A375 or MDA231 and the resulting transfectants were tested for stimulation of TIL1200. DNA sequences of the plasmid clones were determined with an automated DNA sequencer (Model 373A; Applied Biosystems, Inc.), using the Taq DyeDeoxy terminator cycle sequencing kit (Applied Biosystems, Inc.) using the manufacturer's instructions.

#### *Peptide synthesis and identification of antigenic peptides*

Peptides were synthesized by a solid phase method using a Gilson AMS 422 multiple peptide synthesizer. The peptides were purified by HPLC on a Vydac C-4 column with 0.05% TFA/water-acetonitrile. To identify antigenic peptides, TIL lysis of T2 RET-cells preincubated with peptides for 2 hour (h) was measured using a  $^{51}\text{Cr}$  release cytotoxicity assay.

#### *Treatment of a patient with metastatic melanoma using TIL 1200*

A 29 year old male patient, designated patient number 1200, with a widely metastatic melanoma who had previously failed chemotherapy and radiation therapy was treated with a single preparatory dose of 25 mg/Kg cyclophosphamide followed by the intravenous infusion of  $1.6 \times 10^{11}$  TIL (including  $9.1 \times 10^9$  Indium-111 labeled TIL) plus 7 doses of IL-2 at 720,000 IU/Kg given every 8 hours. A second cycle of treatment with TIL and IL-2 was given three weeks later. Radionuclide scans showed localization of TIL in tumor deposits (Figure 3A). Biopsy of subcutaneous tumors on days 8 and 11 after treatment showed significant localization of TIL to tumor (ratios of injectate per gram in tumor compared to normal tissue were 14.9 and 14.0 respectively). The patient's cancer regressed rapidly following the first course of treatment. By three months

after treatment 2 of 3 liver lesions had disappeared and a third lesion shrank by 50%. Multiple subcutaneous metastases regressed completely as shown in Figure 3B (the product of perpendicular diameters of individual lesions are shown).

5

*Characterization of in vitro function of TIL1200*

A number of TIL lines established from HLA-A2<sup>+</sup> melanoma patients lysed melanoma cell lines in a class I MHC-restricted fashion (Kawakami, Y., et al. (1992) J. Immunol. 148; 638-643), and were shown to release IFN $\gamma$ , tumor necrosis factor-alpha (TNF $\alpha$ ) or granulocyte-macrophage colony stimulating factor (GM-CSF) when cocultured with the same tumor cell lines (Hom, S.S., et al. (1993) J. Immunother. 13; 18-30). A CD8<sup>+</sup> CTL line, TIL1200, established from a metastatic subcutaneous tumor mass of patient 1200, lysed fresh autologous melanoma cells as well as 10 of 15 HLA-A2<sup>+</sup> allogeneic melanoma cell lines, but did not lyse 16 of 18 HLA-A2<sup>-</sup> melanoma cell lines or 6 of 8 HLA-A2<sup>+</sup> non-melanoma cell lines (Shilyansky, J., et al. (1993) Proc. Natl. Acad. Sci. USA, 91, 2829-2833, unpublished data). Table 7 shows a cytotoxicity assay against 5 representative HLA-A2<sup>+</sup> melanoma cell lines that were lysed by TIL1200, 4 representative HLA-A2<sup>+</sup> melanoma cell lines that were not lysed by TIL1200, and one HLA-A2<sup>-</sup> melanoma cell line. TIL1200 also secreted IFN- $\gamma$  when cocultured with HLA-A2<sup>+</sup> normal cultured melanocytes established from neonatal foreskin as well as HLA-A2<sup>+</sup> melanoma cell lines (Table 8). Therefore, TIL1200 appeared to recognize a non-mutated self peptide expressed in most melanomas and cultured neonatal melanocytes in an HLA-A2 restricted fashion.

35

*Cloning of the cDNA coding for a melanoma antigen recognized by T cells*

A cDNA library in  $\lambda$  pCEV27 from the HLA-A2<sup>+</sup> 501mel melanoma cell line, which was lysed by most HLA-A2

° restricted melanoma specific TIL, was stably transfected  
into the highly transfectable HLA-A2<sup>+</sup> melanoma antigen  
negative MDA231 clone 7 or A375 clone 1-4. G418 resistant  
cells were selected and approximately 6700 individual  
transfectants from each cell line were isolated and  
5 screened based on their ability to stimulate IFN- $\gamma$   
secretion from TIL1200. Six DNA fragments were isolated  
by PCR using SP6/T7 primers flanking the integrated DNA  
from four MDA231 and one A375 transfectants that were  
positive in a second screening and were cloned into the  
10 mammalian expression vector pcDNA3 (Invitrogen).

These fragments in the pcDNA3 vector were transiently  
expressed in the COS7 cells with or without pcDNA3-HLA-  
A2.1. Transfection into COS7 of one of the cDNAs tested,  
cDNA25, along with HLA-A2.1 reproducibly conferred the  
15 ability to stimulate secretion of IFN- $\gamma$  from TIL1200. The  
stable transfection of cDNA25 into A375 also stimulated  
IFN- $\gamma$  release from TIL1200 (Table 9, Exp.1 and Exp. 2). A  
2.2 Kb band detected by Northern blot analysis of the  
melanoma using the cDNA25 probe suggested that the cloned  
20 1.6Kb fragment was not a full length cDNA. Comparison  
with the GenBank database of the consensus DNA sequence of

25

30

35

Table 7. Specificity of Antigen Recognition by TIL1200:  
Lysis of HLA-A2<sup>+</sup>, gp100<sup>+</sup> Melanoma Cell Lines

Target HLA-A2    gp100    TIL1200    LAK  
FACS Northern

5

10

15

				(% specific lysis)	
501mel	+	+	+	46	78
526mel	+	+	+	39	74
624mel	+	+	+	33	76
952mel	+	+	+	25	76
Malme3M	+	+	+	43	70
C32	+	-	-/+*	6	82
RPMI7951	+	-	-	9	67
WM115	+	-	-	5	68
HS695T	+	-	-	3	87
397mel	-	+	+	0	70

5 hour (h) <sup>51</sup>Cr release assay was performed to measure cellular cytotoxicity at an effector: target ratio of 40:1 as previously described (Kawakami, Y. et al. (1988) J. Exp. Med. 168:2183-2191). The expression of HLA-A2 and gp100 recognized by monoclonal antibody HMB45 (Enzo Diagnostics, New York, NY) was measured by flow cytometry (FACS). The expression of gp100 RNA was analyzed by Northern blot with a cDNA25 probe.

\*-/+ indicates very weak positive.

30

35

Table 8. Specificity of Antigen Recognition by TIL1200:  
Recognition of HLA-a-a2<sup>+</sup> Neonatal Melanocytes

	Stimulator HLA-A2		TIL1200	TIL888
			(pgIFN- $\gamma$ /ml)	
5	501mel	+	562	0
	624mel	+	439	0
	397mel	-	0	0
	888mel	-	0	1970
10	NHEM493	-	441	0
	NHEM527	+	418	0
	NHEM530	+	164	0
	NHEM616	+	53	0
	FM725	+	107	0
15	FM801	+	250	343
	NHEM483	-	0	0
	NHEM680	-	0	0
	HA002	-	0	0
-----				

The IFN- $\gamma$  secretion by TIL was measured by ELISA as previously described in Example 1. The amount of IFN- $\gamma$  secreted by TIL alone was subtracted (88 pg/ml for TIL888 and none for TIL 1200). TIL888 is a class I MHC restricted melanoma specific CTL, not restricted by HLA-A2. NHEM, FM, and HA refer to normal cultured melanocyte cell lines, all others are melanoma cell lines.

**Table 9.** Transfection of cDNA 25 into A375 and COS7

	Stimulator cells	Transfected genes	HLA-A2	Secretion by TIL1200 (pgIFN $\gamma$ /ml)
<u>Exp. 1</u>				
5	501mel	none	+	987
	397mel	none	-	0
	A375	none	+	0
10	A375	pcDNA3-25	+	230
<u>Exp. 2</u>				
	501mel	none	+	662
	397mel	none	-	0
15	COS7	none	-	0
	COS7	HLA-A2.1	+	0
	COS7	pcDNA3-25	-	0
20	COS7	HLA-A2.1+pcDNA3-25	+	310
<u>Exp. 3</u>				
	501mel	none	+	908
	397mel	none	-	0
25	COS7	none	-	0
	COS7	HLA-A2.1	+	0
	COS7	pCEV27-FL25	-	0
	COS7	HLA-A2.1+pCEV27-FL25	+	742
30	COS7	pcDNA3-FL25	-	0
	COS7	HLA-A2.1+pcDNA3-FL25	+	801

TIL 1200 secreted IFN- $\gamma$  when coincubated with HLA-A2<sup>+</sup> A375 stably transfected with pcDNA3 containing truncated cDNA25 (pcDNA3-25) (Exp.1) or COS7 transiently transfected with either pcDNA3-25 (Exp. 2), pcDNA3 containing full length cDNA25 (pcDNA3-FL25) or pCEV27 containing full length cDNA25 (pCEV27-FL25) (Exp. 3) along with pcDNA3 containing HLA-A2.1 (HLA-A2.1). HLA-A2 expression was determined by flow cytometry and interferon-gamma secretion was measured by ELISA.

35

3 cDNA25 clones that were independently amplified by PCR revealed that cDNA25 was distinct from two previously registered genes, a gp100 (GenBank Access No. M77348) and Pmel17 (Kwon, B.S., et al. (1991) Proc. Natl. Acad. Sci., USA 88, 9228-9232). The cDNA 25 differed from the gp100 in GenBank (Accession No. M77348, also known as gp95) by two nucleotides, from the Pmel 17 sequence (Kwon et al. (1991) Proc. Natl. Acad. Sciences (USA) 88: 9228-9232) by 3 bases and a 21 base pair deletion. (Figure 5B).

The full length cDNA25 (FL25) was isolated in two plasmids, pCEV27-FL25 or pCDNA3-FL25. Transfection of either plasmid into COS7 along with pCDNA3-HLA-A2.1 conferred to COS7 the ability to induce IFN- $\gamma$  secretion by TIL1200. The amount of IFN- $\gamma$  secretion stimulated by COS7 transfected with the full length DNA plus HLA-A2.1 was similar to that stimulated by 501mel and was higher than that stimulated by COS7 transfected with the truncated cDNA25 possibly due to improved translation starting at the normal AUG initiation codon (Table 9, Exp.2 and 3). Alternatively, the 5' region missing from the truncated cDNA25 may contain other epitopes recognized by clones in TIL1200. The requirement for HLA-A2.1 expression for IFN- $\gamma$  release from TIL1200 and the fact that transfected cells did not stimulate IFN- $\gamma$  secretion from irrelevant TIL (data not shown) demonstrated that the cDNA25 encoded an antigen recognized by TIL1200 in the context of HLA-A2.1 and did not encode a molecule that non-specifically induced IFN- $\gamma$  release from T cells.

The nucleotide and corresponding amino acid sequences of the truncated cDNA 25 and the full length cDNA25 cloned from the 501mel cDNA library by screening with the cDNA25 probe (Figure 5A) were compared with the GenBank sequences of Pmel17 isolated from normal melanocytes and gp100 isolated from the melanoma cell line MEL-1. (Figure 5B). The full length cDNA25 differed from the gp100 amino acid sequence at position 162. This amino



acid difference is possibly caused by polymorphism or mutation in the tumor. cDNA25 had 2 amino acid differences at positions 162 and 274, compared to Pmel17 and did not contain 7 amino acids that existed in Pmel17 at positions 588-594. The amino acid sequence of the truncated cDNA25 that was isolated from the original MDA231 transfectant has a different sequence at the 3' end (from position 649 to the end) due to a frame shift caused by one extra cytidylic acid. It is not clear whether this difference was due to a true allelic difference or to a mutation that occurred during manipulation of the DNA. Nevertheless, TIL1200 appeared to recognize non-mutated peptides located between position 236 and 648. cDNA25 also had 87% similarity in amino acid sequence to cDNA RPE1 (Kim, R., and Wistow, G.J. (1992) Exp. Eye Res. 55: 657-662) specifically expressed in bovine retinal pigment epithelium and 60% similarity to cDNA MMP115 that encoded a melanosomal matrix protein isolated from chicken pigmented epithelial cells (Shilyansky, J., et al. (1993) Proc. Natl. Acad. Sci. USA, 91, 2829-2833).

A gp100 protein was known to be recognized by monoclonal antibody HMB45 (Adema et al., (1993) Am. J. Pathology, 143: 1579-1585). COS7 cells transfected with the full length cDNA25 were evaluated by flow cytometry using this monoclonal antibody. After transient expression of either pCEV27-FL25 or pcDNA3-FL25, COS7 expressed the antigen detected by HMB45 (data not shown).

#### *Expression of RNA for cDNA25*

Northern blot analysis was performed with the cDNA25 probe to evaluate the tissue specific expression of this gene. Ten of 15 melanoma cell lines and 6 of 6 melanocyte cell lines were positive for cDNA 25. Of many normal tissues tested only retina was positive (Figure 6). Seven cell lines from T-cell (TILA, B), B-cells (501EBVB, 836EBVB) and fibroblast (M1) and 20 non-melanoma tumor

cell lines (colon cancer, Collo, SW480, WiDr; breast cancer, MDA231, MCF7, HS578, ZR75; neuroblastoma, SK-N-AS, SK-N-SH; Ewing sarcoma, TC75, RD-ES, 6647; sarcoma 143B; glioma, U138MG, HS683; renal cell cancer, UOK108, UOK117, small cell lung cancer, H1092; Burkitt's lymphoma, Daudi; myeloma HMY) were all negative for cDNA25 (data not shown). Therefore, this gene appeared to be specifically expressed in melanocyte lineage cells, consistent with the expression pattern of previously isolated forms of gp100 when analyzed using monoclonal antibodies, HMB45, NKI/betab, or HMB-50 (Adema, G.J., et al. (1993) Am J Pathology 143: 1579-1585; Gown, A.M., et al., (1986) Am J Pathol 123:195-203; Colombari, R., et al. (1988) Virchows Archiv A Pathol Anat. 413:17-24); Vennegoor, C., et al. (1988) Am. J. Pathol. 130:179-192; Vogel, A.M., and Esclamado R.M. (1988) Cancer Res. 48:1286-1294). The levels of expression of the RNA detected by the cDNA25 probe in cultured neonatal melanocyte cell lines was significantly lower than that in melanoma cell lines. There was a perfect correlation between gp100 expression detected by Northern blot analysis with cDNA25 and flow cytometry using HMB45 antibody and melanoma lysis by TIL1200 in the 10 HLA-A2<sup>+</sup> melanoma cell lines as shown in Table 7.

#### *Identification of the epitope in gp100*

Based on a comparison of the amino acid sequence of the truncated form of cDNA 25 to known binding motifs of HLA-A2.1 (Falk, K., et al. (1992) Nature 351:290-296; Hunt, D.F., et al. (1992) Science 255:1261-1263; Ruppert, J., et al. (1993) Cell 74:929-9937,) 30 peptides of 9 or 10 amino acids in length from cDNA25 were synthesized. TIL1200 lysed the HLA-A2<sup>+</sup> cell line, T2, only when incubated with the peptide LLDGTATLRL (SEQ ID NO: 27 residues 457-486. Figure 5A; ~~SEQ ID NO: 33~~) but not when incubated with the other 29 peptides (Table 10, Figure 5A). Only peptide LLDGTATLRL (~~SEQ ID NO: 33~~) was able to

also stimulate IFN- $\gamma$  secretion by TIL 1200 (data not shown).

Many melanoma-specific CTL derived from TIL appear to recognize non-mutated self peptides derived from melanocyte-melanoma lineage specific proteins, since these TIL recognize most melanoma cell lines and normal cultured melanocytes sharing the appropriate restriction element. (Anichini, A., et al. (1993) J. Exp. Med. 177:989-998; Kawakami, Y., et al. (1993) J. Immunother. 14:88-93). In an attempt to isolate and identify melanoma antigens of value in the immunotherapy of melanoma patients, TIL, TIL1200 were used that, when transferred into a patient with metastatic cancer, localized to the tumor site and was associated with a dramatic tumor regression. It has been shown that, in contrast to non-activated lymphocytes and lymphokine activated killer cells, autologous TIL localize to tumor sites. This localization correlated with the ability of these TIL to mediate tumor regression (data not shown). TIL1200 which was a TIL line containing multiple CTL species recognized a tumor antigen in the context of HLA-A2, which is the most frequently expressed class I MHC antigen (about 50% of individuals) and has been shown to be a dominant restriction element for the induction of melanoma specific CTL. (Crowley, N.J., et al. (1991) J. Immunol. 146, 1692-1699).

By cDNA expression cloning using T cell recognition for screening, a cDNA (Figure 4; SEQ ID NO: 26) encoding an antigen recognized by TIL1200 and identified as a form of gp100, a membrane glycoprotein also recognized by monoclonal antibodies, HMB45, HMB50 or NKI/betab has been identified. (Adema, G.J., et al. (1993) Am J Pathology 143, 1579-1585. Gown, A.M., et al. (1986) Am J Pathol 123, 195-203. Colombari, R., et al. (1988) Virchows Archiv A Pathol Anat. 413, 17-24; Vennegoor, C., et al. (1988) Am. J. Pathol. 130, 179-192; Vogel, A.M., and Esclamado R.M. (1988) Cancer Res. 48,

1286-1294). These antibodies are highly specific for melanocyte lineage tissues and strongly stain most melanoma cells. NKI/betab also reacts with adult melanocytes in normal skin (Vennegoor, C., et al. (1988)

5

10

15

20

25

30

35

Table 10. TIL1200 Lysis of the T2 HLA-A2<sup>+</sup> Cell Line  
Pulsed with the Peptide, LLDGTATLRL

Target	HLA-A2	Peptide*	TIL1200	TIL1235 <sup>+</sup>
		(ug/ml)	(% specific lysis)**	
501mel	+	0	66	51
397mel	-	0	1	0
T2	+	0	2	1
T2	+	40	28	ND <sup>++</sup>
T2	+	10	32	0
T2	+	1	24	ND
T2	+	0.1	6	ND
T2	+	0.01	0	ND
T2	+	0.001	2	ND

15 \*TIL1200 lysed T2 cells pulsed with the 10-mer peptide, LLDGTATLRL(457-466), but not other 29 peptides SEQ ID NO: 27 (residues 273-281, 297-306, 373-381, 399-407, 399-408, 409-418, 456-464, 463-471, 465-473, 476-485, 511-520, 519-528, 544-552, 544-553, 570-579, 576-584, 576-585, 585-593, 592-600, 597-605, 597-606, 602-610, 602-611, 603-611, 605-614, 606-614, 606-615, 619-627, 629-638)

20 \*TIL1235 is an HLA-A2 restricted melanoma specific CTL that does not recognize gp100.

\*\*E:T of 50:1

25 ++ND, not done.

30

35

Am. J. Pathol. 130, 179-192). Immunoelectron-microscopic studies using either HMB45 or NKI/betab antibody revealed that a gp100 protein was mainly located in a membrane and filamentous matrix of stage I and II melanosomes in the cytoplasm (Vennegoor, C., et al. (1988) Am. J. Pathol. 130, 179-192; Schaumburg-Lever, G., et al. (1991) J. Cutan. Pathol. 18, 432-435). By a completely independent procedure, a cDNA encoding another form of gp100 was also isolated by screening with a rabbit polyclonal antiserum against gp100 (Adema, G.J., et al. (1993) Am J Pathology 143:1579-1585) and TIL1200 also lysed HLA-A2<sup>+</sup> cell lines transfected with this cDNA clone (Bakker, A.B.H. et al. (1994) J. Exp. Med. 179:1005-1009).

The existence of T cells reactive to the self-antigen gp100 in tumors and the possible enrichment of these T cells at the tumor site as a possible consequence of the specific accumulation and expansion of antigen reactive cells (Sensi, M. et al., (1993) J. Exp. Med 178:1231-1246) raises important questions about the nature of the immune response to self antigens on growing cancers and about the mechanisms of immunologic tolerance to self-antigens. The increased expression of gp100 on melanoma cells relative to that in melanocytes demonstrated by Northern blot analysis or the unique inflammatory conditions that might exist at the tumor site, which may be associated with the secretion of cytokines and expression of costimulatory molecules on the cell surface, could break tolerance to gp100. Depigmentation has been reported to be associated with a good prognosis (Nordlund, J.J., et al. (1983) J. Am. Acad. Dermatol. 9:689-695; Bystry, J-C, et al. (1987) Arch. Dermatol., 123:1053-1055) and with clinical response to chemoimmunotherapy (Richards, J.M., et al. (1992) J. Clin. Oncol. 10:1338-1343) in melanoma patients. Sporadic vitiligo in has been seen patients receiving melanoma-specific TIL but adverse ophthalmologic effects that might be related to melanocyte

destruction has not been observed. Patient 1200 did not develop vitiligo or any ophthalmologic side-effects.

The gp100 protein (Figure 5A; SEQ ID NO: 27) and the ten amino acid peptide identified may represent a human tumor rejection antigen since the transfer into patient 1200 of TIL1200 plus IL2 was associated with cancer regression. The traffic of TIL1200 to tumor deposits in vivo and the rapidity of the antitumor response are characteristics of the response to TIL therapy, although IL2 may also have been involved in the tumor rejection. Adoptive transfer of 3 other TIL lines which recognized gp100 as well as MART-1 also mediated tumor regression (data not shown).

Tyrosinase (Brichard, V., et al. (1993) J. Exp. Med. 178, 489-495) and MART-1 (see Example 1) have been identified as melanoma antigens recognized by HLA-A2 restricted CTL. Another antigen, MAGE-1 is recognized by HLA-A1 restricted melanoma-specific CTL and is expressed on a variety of cancer cells as well as testis (Van Der Bruggen, P. et al. (1991) Science, 254:1643-1647). However, none of the ten HLA-A2 restricted TIL recently developed appeared to recognize MAGE-1 (Zakut, R., et al. (1993) Cancer Res. 53: 5-8).

The wide expression of gp100 proteins in melanomas, the recognition of a peptide by T cells infiltrating into tumor, its restriction by HLA-A2, present in 50% of individuals, and the association of anti gp100 reactivity with cancer regression in patient 1200 imply that the gp100 antigen in particular the novel immunogenic peptides derived from the gp100 amino acid sequence (Figure 5A; SEQ ID NO: 27) may be particularly useful for the development of active immunotherapies for patients with melanoma.

#### Example 4

MART-1 Vaccines As A Treatment For Melanoma In Mammals

MART-1 vaccines may be efficacious in treating mammals afflicted with melanoma. For example, MART-1 vaccines may be administered to individuals. Mammals can be immunized with the recombinant proteins described  
5 herein in ranges of 1mg-100mg. Alternatively mammals, preferably humans may be immunized with the MART-1 nucleic acid sequence inserted into a viral vector such as vaccinia virus, adenovirus or fowl pox virus. A range of about  $10^6$ - $10^{11}$  viral particles carrying the MART-1 nucleic  
10 acid sequences can be administered per mammal, preferably a human. The mammals will be monitored for antibodies to the immunogen or increase in cytotoxic lymphocytes (CTL) recognizing the immunogen by conventional methods or alleviation of clinical signs and symptoms of the active  
15 disease. Specific parameters to be assessed include production of immune cells that recognize the vaccine antigen or tumor regression. Such vaccines may be administered either prophylactically or therapeutically. Mammals may also be immunized with the MART-1 nucleic acid  
20 sequence inserted into a retroviral vector. Suggested dose ranges of the antigen in retroviruses are  $10^6$ - $10^{11}$  viral particles per mammal, preferably a human. Response and efficacy of the retroviral vaccines will be assessed as described above.

#### Example 5

Use Of Lymphocytes Sensitized To  
Immunogenic Peptides Derived From  
Melanoma Antigens For Therapeutically  
Treating Mammals Afflicted With  
30 Melanoma

T-lymphocytes presensitized to the melanoma antigen may be effective in therapeutically treating mammals afflicted with melanoma. The T-lymphocytes will  
35 be isolated from peripheral blood lymphocytes or tumor infiltrating lymphocytes and exposed in vitro to the



MART-1 protein or peptide. T-lymphocytes are isolated from peripheral blood or melanoma tumor suspensions and cultured in vitro (Kawakami, Y. et al. (1988) J. Exp. Med. 168: 2183-2191). The T-lymphocytes are exposed to the MART-1 peptide AAGIGILTV for a period of about 1-16 hours at a concentration of 1 to 10mg/ml. T-lymphocytes exposed to the antigen will be administered to the mammal, preferably a human at about  $10^9$ - $10^{12}$  lymphocytes per mammal. The lymphocytes may be administered either intravenously, intraperitoneally or intralesionally. This treatment may be administered concurrently with other therapeutic treatments such as cytokines, radiotherapy, surgical excision of melanoma lesions and chemotherapeutic drugs, adoptive T lymphocyte therapy.

The present invention is not to be limited in scope by the nucleic acid sequences deposited, since the deposited embodiments is intended as a single illustration of one aspect of the invention and any sequences which are functionally equivalent are within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the dependent claims. It is also to be understood that all base pair sizes given for nucleotides are approximated and are used for purpose of description.

82